



Belle: recent results and future plans

Peter Križan

University of Ljubljana and J. Stefan Institute



Contents

Experimental apparatus: Belle at KEK-B

CP violation in the B system

Searching for New Physics: FCNC processes

- Observation of $b \rightarrow d$ penguins: $B \rightarrow \rho\gamma, \omega\gamma$ decays
- CP violation in $b \rightarrow s$ decays
- A_{fb} vs q^2 in $B \rightarrow K^* l^+ l^-$ decays

Plans for the future: a Super B factory



Belle Collaboration



13 countries, 55 institutes, ~400 collaborators


January 5, 2006

University of Dortmund

Peter Krizan, Ljubljana



Belle spectrometer at KEK-B




January 5, 2006

University of Dortmund

Peter Krizan, Ljubljana



The KEKB Collider





January 5, 2006

University of Dortmund



Normal injection

Continuous injection



661/pb/day

→1182/pb/day

January 5, 2006


University of Dortmund

Peter Krizan, Ljubljana



Belle/KEKB Luminosity Milestone: $500 \text{ fb}^{-1} = 0.5 \text{ ab}^{-1}$

Accumulated > 500 M BB-pairs



1 fb⁻¹/day

Total = 512 fb⁻¹

Today: some results with 350 fb⁻¹ (386 x 10⁶) B B pairs

as well as results based on 253 fb⁻¹ (275 x 10⁶) B B pairs

Peter Krizan, Ljubljana




B factory physics program

B factory main task: measure CP violation in the system of B mesons

specifically: various measurements of complex elements of **Cabbibo-Kobayashi-Maskawa matrix**

CKM matrix is unitary

deviations could signal processes not included in SM



$$\begin{pmatrix} V_{ud} & V_{us} & V_{ub} \\ V_{cd} & V_{cs} & V_{cb} \\ V_{td} & V_{ts} & V_{tb} \end{pmatrix} = \begin{pmatrix} 1-\lambda^2/2 & \lambda & A\lambda^3(\bar{\rho}-i\bar{\eta}) \\ -\lambda & 1-\lambda^2/2 & A\lambda^2 \\ A\lambda^3(1-\bar{\rho}-i\bar{\eta}) & -A\lambda^2 & 1 \end{pmatrix}$$


January 5, 2006

University of Dortmund

Peter Krizan, Ljubljana



Three Angles: (ϕ_1, ϕ_2, ϕ_3) or (β, α, γ)



Big Questions: *Are determinations of angles consistent with determinations of the sides of the triangle? Are angle determinations from **loop** and **tree** decays consistent?*



January 5, 2006

University of Dortmund

Peter Krizan, Ljubljana



Mixing Induced CP Violation



$$\rightarrow A_{CP}(t) = \frac{\Gamma(\bar{B}^0(t) \rightarrow f_{CP}) - \Gamma(B^0(t) \rightarrow f_{CP})}{\Gamma(\bar{B}^0(t) \rightarrow f_{CP}) + \Gamma(B^0(t) \rightarrow f_{CP})} = \xi_f \sin 2\phi \sin \Delta m_B t$$

$\xi_f = \pm 1$ for $CP = \pm 1$


January 5, 2006

University of Dortmund

Peter Krizan, Ljubljana



Principle of CPV Measurement



January 5, 2006

University of Dortmund

Peter Krizan, Ljubljana



If there is **more than one diagram** and additional weak phases, there is the possibility of **direct CPV** and a new term with a **$\cos(\Delta mt)$** time dependence.

$$P(B \rightarrow f_{CP}; t) = \frac{e^{-t/\tau_B}}{4\tau_B} [1 + q \cdot \{A \cos(\Delta mt) + S \sin(\Delta mt)\}]$$

with $q = \pm 1$

If integrated over all times $(-\infty, +\infty)$, the asymmetry with the **$\sin(\Delta mt)$** term **vanishes**, while the term with **$\cos(\Delta mt)$** **remains**.



January 5, 2006

University of Dortmund


Peter Krizan, Ljubljana



Reconstruction of B meson decays



Reconstructing rare B meson decays at Y(4s): use two variables, **beam constrained mass M_{bc}** and **energy difference ΔE**



$$\Delta E \equiv \sum E_i - E_{CM}/2$$


$$M_{bc} = \sqrt{(E_{CM}/2)^2 - (\sum \vec{p}_i)^2}$$

Peter Krizan, Ljubljana




2005: $B^0 \rightarrow J/\psi \bar{K}^0$ with 386 M $B\bar{B}$ pairs

$B^0 \rightarrow J/\psi K_S^0$



$$M_{bc} = \sqrt{E_{beam}^{*2} - P_{J/\psi K_S}^{*2}}$$

$B^0 \rightarrow J/\psi K_L^0$





p_B^* (momentum in CM)

January 5, 2006

University of Dortmund

Peter Krizan, Ljubljana





Evidence and Observation of Direct CP Violation in B Decays

DCPV in $B^0 \rightarrow \pi^+ \pi^-$ and $B^0 \rightarrow K^- \pi^+$,

hep-ex/0502035 (PRL 95, 101801(2005)); hep-ex/0507045

Asymmetries in the Dalitz plot of $B^\pm \rightarrow K^\pm \pi^+ \pi^-$

hep-ex/0509001

January 5, 2006

University of Dortmund

Peter Krizan, Ljubljana



Importance of direct CPV in B decays

"The final, completely definitive death of any superweak theory will come from the observation of direct CP violation in the B system....."

Evidence for such direct CP violation would be given by the difference between the asymmetry parameters in a decay such as $B \rightarrow \pi^+ \pi^-$ from that of $B \rightarrow J/\psi K_S$. This can be considered the **ϵ' experiment for the B system.**"

Lincoln Wolfenstein, 1999

January 5, 2006

University of Dortmund

Peter Krizan, Ljubljana



Direct CPV asymmetry in B Decays

Asymmetry in B decay rates

$$A_{dir} \equiv \frac{\Gamma(\bar{B} \rightarrow \bar{f}) - \Gamma(B \rightarrow f)}{\Gamma(\bar{B} \rightarrow \bar{f}) + \Gamma(B \rightarrow f)}$$

$$= \frac{2r \sin \phi \sin \delta}{1 + r^2 + 2r \cos \phi \cos \delta}$$

$r = |P|/|T|$, $\phi = \text{weak phase diff}$
 $\delta = \text{strong phase diff}$

The direct CP asymmetry (A_{dir}) can be large if two amplitudes have comparable sizes, **different weak phases as well as a strong phase difference**. This can happen in certain B decays due to the interference of penguin (P) and tree (T) decays.


January 5, 2006

University of Dortmund

Peter Krizan, Ljubljana



Hierarchy of diagrams for $B \rightarrow \pi\pi, K\pi$ decays



Possibility of tree-penguin interference.

N.B. in $B \rightarrow \pi\pi$ the two diagrams are the same order in λ

January 5, 2006

University of Dortmund

Peter Krizan, Ljubljana



Experimental Situation for $B \rightarrow \pi^+ \pi^-$ in 2004



Belle 152 M $\overline{B}B$

with 372 ± 32 $B^0 \rightarrow \pi^+ \pi^-$ events

$$S_{\pi\pi} = -1.00 \pm 0.21 \pm 0.07$$

$$A_{\pi\pi} = +0.58 \pm 0.15 \pm 0.07$$

PRL 93, 021601 (2004)

**5.2 σ CPV,
First evidence for DCPV (3.2 σ)**



BABAR 227M $\overline{B}B$

with 467 ± 33 $B^0 \rightarrow \pi^+ \pi^-$ events

$$S_{\pi\pi} = -0.30 \pm 0.17 \pm 0.03$$

$$A_{\pi\pi} = +0.09 \pm 0.15 \pm 0.04$$

hep-ex/0501071, to appear in PRL

Also $\sim 3.2s$ discrepancy between Belle and BaBar


January 5, 2006

University of Dortmund

Peter Krizan, Ljubljana



$B \rightarrow \pi^+ \pi^-$ time evolution



666 ± 43 $B \rightarrow \pi^+ \pi^-$ signal events

ΔE -Mbc 2D fits to individual time intervals

Bkg subtracted fit projections for $B \rightarrow \pi^+ \pi^-$

2005 sample

January 5, 2006

University of Dortmund

Peter Krizan, Ljubljana



Summary of Belle $B^0 \rightarrow \pi^+ \pi^-$ CPV results


$$A_{\pi\pi} = +0.56 \pm 0.12 \pm 0.06 \quad \begin{array}{l} \text{1st error statistical,} \\ \text{2nd systematic} \end{array}$$
$$S_{\pi\pi} = -0.67 \pm 0.16 \pm 0.06$$

- Compelling evidence for direct CP violation in $B \rightarrow \pi^+ \pi^-$ with 4.0σ significance
- Confirms previous Belle results.
- Isospin analysis for this mode alone gives (95.4% C.L) $0^\circ < \phi_2 < 19^\circ$ & $71^\circ < \phi_2 < 180^\circ$





Consistency Checks with time-integrated fits

$$A_{\pi\pi} = +0.52 \pm 0.14 \quad \text{Counting experiment consistent with unbinned time-dependent fits.}$$




Visible indication of direct CP violation.






2005: "Observation" of Direct CPV in $B \rightarrow K^-\pi^+$

Belle update with
 386×10^6 B Bbar pairs
(hep-ex/0507045)



One more nail in the
Superweak coffin.



Significance
 5.0σ

$$A_{CP}(K^+\pi^-) \equiv \frac{N(\bar{B} \rightarrow K^-\pi^+) - N(B \rightarrow K^+\pi^-)}{N(\bar{B} \rightarrow K^-\pi^+) + N(B \rightarrow K^+\pi^-)} = 0.113 \pm 0.022 \pm 0.008.$$

January 5, 2006

University of Dortmund

Peter Krizan, Ljubljana



Interpretation: Direct CP violation+SU(3)

The results support the expectation from SU(3) symmetry that

$$A_{CP}(K^+\pi^-) \sim -\frac{1}{3} A_{CP}(\pi^+\pi^-)$$

N.G. Deshpande and X.-G. He, PRL 75, 1703 (1995)

M. Gronau and J.L. Rosner, PLB 595, 339 (2004)

$$A_{CP}(K^+\pi^-) = -0.115 \pm 0.018$$

HFAG summer 2005

$$-\frac{1}{3} A_{CP}(\pi^+\pi^-) = -0.19 \pm 0.04$$

Belle measurement

January 5, 2006

University of Dortmund

Peter Krizan, Ljubljana




A new approach to direct CPV using the Dalitz plot in $B^\pm \rightarrow K^\pm \pi^+ \pi^-$ (hep-ex/0509001)

Sample used for $B^\pm \rightarrow K^\pm \pi^+ \pi^-$ study:


contains $2248 \pm 79 B^-$, $2038 \pm 76 B^+$

Fix the resonant substructure, then allow both the phase and amplitude to be different for B^+ and B^- decays.

For each resonant amplitude replace $ae^{i\delta}$ with $ae^{i\delta}(1 \pm b e^{i\phi})$



Combined Dalitz plot, signal region



Jar FIG. 9: Illustration of the amplitude parametrization with Eq. 7.

Peter Krizan, Ljubljana



Evidence for CP Violation in the Decay $B^\pm \rightarrow \rho^0 K^\pm$

TABLE I: Results of the best fit to $K^\pm \pi^\pm \pi^\mp$ events in the B signal region. The first quoted error is statistical and the second is the model dependent uncertainty. The quoted A_{CP} significance is statistical only.

Channel	Fraction (%)	δ ($^\circ$)	b	φ ($^\circ$)	A_{CP} significance (σ)
$K^*(892)\pi^\pm$	$13.0 \pm 0.8^{+0.5}_{-0.7}$	0 (fixed)	$0.078 \pm 0.033^{+0.012}_{-0.003}$	$-18 \pm 44^{+5}_{-13}$	2.6
$K_0^*(1430)\pi^\pm$	$65.5 \pm 1.5^{+2.2}_{-3.9}$	$55 \pm 4^{+1}_{-5}$	$0.069 \pm 0.031^{+0.010}_{-0.008}$	$-123 \pm 16^{+4}_{-5}$	2.7
$\rho(770)^0 K^\pm$	$7.85 \pm 0.93^{+0.64}_{-0.59}$	$-21 \pm 14^{+14}_{-19}$	$0.28 \pm 0.11^{+0.07}_{-0.09}$	$-125 \pm 32^{+10}_{-85}$	3.9
$\omega(782)K^\pm$	$0.15 \pm 0.12^{+0.03}_{-0.02}$	$100 \pm 31^{+38}_{-21}$	0 (fixed)	-	-
$f_0(980)K^\pm$	$17.7 \pm 1.6^{+1.1}_{-3.3}$	$67 \pm 11^{+10}_{-11}$	$0.30 \pm 0.19^{+0.05}_{-0.10}$	$-82 \pm 8^{+2}_{-4}$	1.6
$f_2(1270)K^\pm$	$1.52 \pm 0.35^{+0.22}_{-0.37}$	$140 \pm 11^{+18}_{-7}$	$0.37 \pm 0.17^{+0.11}_{-0.04}$	$-24 \pm 29^{+14}_{-20}$	2.7
$f_X(1300)K^\pm$	$4.14 \pm 0.81^{+0.31}_{-0.30}$	$-141 \pm 10^{+8}_{-9}$	$0.12 \pm 0.17^{+0.04}_{-0.07}$	$-77 \pm 56^{+88}_{-43}$	1.0
Non-Res.	$34.0 \pm 2.2^{+2.1}_{-1.8}$	$\delta_1^{nr} = -11 \pm 5^{+3}_{-3}$ $\delta_2^{nr} = 185 \pm 20^{+62}_{-19}$	0 (fixed)	-	-
$\chi_{c0} K^\pm$	$1.12 \pm 0.12^{+0.24}_{-0.08}$	$-118 \pm 24^{+37}_{-38}$	$0.15 \pm 0.35^{+0.08}_{-0.07}$	$-77 \pm 94^{+154}_{-11}$	0.7

$$A_{CP}(B^\pm \rightarrow \rho^0 K^\pm) = 0.28 \pm 0.10^{+0.07}_{-0.09} \quad (3.9\sigma)$$

Significance varies from 3.7σ to 4.0σ depending on the model for the resonant substructure (add or remove modes, change nr model, cpv in $b \rightarrow u$ background).

January 5, 2006

University of Dortmund

Peter Krizan, Ljubljana



Evidence for CP Violation in the Decay $B^\pm \rightarrow \rho^0 K^\pm$

B- vs B+ in the ρ and f^0 (980)
 $m_{\pi\pi}$ region

- helicity
hemisphere

+helicity
hemisphere




FIG. 7: $M(\pi^+\pi^-)$ mass spectra for B^- (left column) and B^+ (right column) for different helicity regions: (a,b) no helicity cuts; (c,d) $\cos\theta_H^+ < 0$; (e,f) $\cos\theta_H^+ > 0$. Points with error bars are data, the open histogram is the fit result and the hatched histogram is the background component.

January 5, 2006



Comparison of result to predictions and BaBar

Belle Data: $A_{CP}(B^\pm \rightarrow \rho^0 K^\pm) = 0.28 \pm 0.10^{+0.07}_{-0.09}$ (3.9 σ)



First evidence for DCPV in a charged meson decay

Cheng, Gronau, Luo, Rosner, Suprun; PRD 69, 034001 (2004)

$$A_{CP}(B^\pm \rightarrow \rho^0 K^\pm) = 0.21 \pm 0.10$$

M. Beneke and M. Neubert; Nucl. Phys. B675, 333 (2003)

$$A_{CP}(B^\pm \rightarrow \rho^0 K^\pm) = -13.6^{+4.5+6.9+3.7+62.7}_{-5.7-4.4-3.1-55.4}\%$$

Four representative scenarios (-27.3, -9.3, 26.6, 31.7)%

BaBar Data: $A_{CP}(B^\pm \rightarrow \rho^0 K^\pm) = 0.32 \pm 0.13 \pm 0.06^{+0.08}_{-0.05}$ (2.4 σ)

January 5, 2006


University of Dortmund

Peter Krizan, Ljubljana



Why FCNC decays?

Flavour changing neutral current (FCNC) processes (like $b \rightarrow s$, $b \rightarrow d$) are forbidden at the tree level in the Standard Model. Proceed only at low rate via higher-order loop diagrams. Ideal place to search for new physics.



January 5, 2006

University of Dortmund

Peter Krizan, Ljubljana




A large number of $b \rightarrow s$ modes are known, where are the $b \rightarrow d$ penguins ?

Suppressed by $|V_{td}/V_{ts}|^2$ vs $b \rightarrow s\gamma$

Interesting:

Measurement of $|V_{td}/V_{ts}|$

CP violation could be sizeable in SM (order 10%)



$$\frac{\mathcal{B}(B \rightarrow (\rho, \omega)\gamma)}{\mathcal{B}(B \rightarrow K^*\gamma)} = S_\rho \left| \frac{V_{td}}{V_{ts}} \right|^2 \left(\frac{1 - m_\rho^2/M_B^2}{1 - m_{K^*}^2/M_B^2} \right)^3 \zeta^2 [1 + \Delta R]$$

Addresses the same physics issue as B_s - B_s mixing (future Tevatron RunII +LHCb goal).

January 5, 2006

University of Dortmund

Peter Krizan, Ljubljana





V_{td}/V_{ts} from $B \rightarrow \rho\gamma, \omega\gamma$

The measured branching fraction, $\mathcal{B}(B \rightarrow (\rho\omega)\gamma) = (1.34^{+0.34}_{-0.31} \text{ } ^{+0.14}_{-0.10}) \times 10^{-6}$, translates to

$$|V_{td}/V_{ts}| = 0.200^{+0.026}_{-0.025}(\text{exp.})^{+0.038}_{-0.029}(\text{theo.}),$$

which is compatible with SM constraints based on fits using measurements of other CKM parameters.



hep-ex/0506079

First Observation of $b \rightarrow d \gamma$

January 5, 2006

University of Dortmund

Peter Krizan, Ljubljana




Implications of Belle's observation of $b \rightarrow d \gamma$

Together with the evidence of $B \rightarrow K^0 K$ modes, Belle has demonstrated the existence of a new quark level transition: $b \rightarrow d$

+ measurement of $|V_{td}/V_{ts}|$

$$\frac{\text{BR}(B \rightarrow (\rho/\omega)\gamma)}{\text{BR}(B \rightarrow K^*\gamma)} \propto \left| \frac{V_{td}}{V_{ts}} \right|^2$$

SU(3) breaking correction
weak annihilation diagram for $\text{BR}(B \rightarrow \rho/\omega\gamma)$




Januar

Ljubljana




How can New Physics contribute to $b \rightarrow s$?


$B^0 \rightarrow \phi K^0$




$B^0 \rightarrow \eta' K^0$



New physics in loops?



Many new phases are possible in SUSY



January 5, 2006

University of Dortmund


Peter Krizan, Ljubljana



Searching for new physics phases in CP violation measurements in $b \rightarrow s$ decays

Example:

no KM phase



SM: $\sin 2\phi_1^{\text{eff}} = \sin 2\phi_1$ from $B \rightarrow J/\psi K^0$ ($b \rightarrow c \bar{c} s$)
 unless there are other, non-SM particles in the loop

January 5, 2006

University of Dortmund

Peter Krizan, Ljubljana





Belle 2005 update:

hep-ex/0507037

$B \rightarrow \phi K^0 : K^0 \rightarrow K_S \text{ or } K_L$ ($386 \times 10^6 B\bar{B}$ pairs)

ϕK_S

ϕK_L



$\sin 2\phi_1 = +0.19 \pm 0.32$

$\sin 2\phi_1 = +1.54 \pm 0.59$

$\sin(2\phi_1)(B \rightarrow \phi K^0) = 0.44 \pm 0.27 \pm 0.05$

January 5, 2006


University of Dortmund

Peter Krizan, Ljubljana



"Compelling Evidence" for CP Violation in a $b \rightarrow s$ mode

$\eta' K^0$ (background subtracted)



$\sin 2\phi_1 = +0.62 \pm 0.12 \pm 0.04$
 $A = -0.04 \pm 0.08 \pm 0.06$

significance $> 4\sigma$

January 5, 2006


University of Dortmund

Peter Krizan, Ljubljana



Many $b \rightarrow s$ modes were studied:

~all are systematically below $\sin(2\beta)$ value from $B \rightarrow J/\psi K^0$ modes




New Physics ?

$\Delta \sin 2\phi_1^{\text{eff}}$ in $b \rightarrow \bar{s}q\bar{q}$ golden modes (July 2005)

Very large effects of order unity, $\Delta S=1$, are now ruled out.

Theory corrections are small and opposite in sign to the measured deviations.






Projection for Super B Factory (50ab⁻¹)

Are there New Physics effects of order 0.2 in b→s CPV ?

Super B Factory level statistics will allow us to answer this question.




January 5, 2006

University of Dortmund

Peter Krizan, Ljubljana



Another FCNC decay: $B \rightarrow K^* l^+ l^-$



$b \rightarrow s l^+ l^-$ was first measured in $B \rightarrow K l^+ l^-$ by Belle (2001).

Important for further searches for the physics beyond SM

$$\frac{d\Gamma(b \rightarrow s l^+ l^-)}{d\hat{s}} = \left(\frac{\alpha_{em}}{4\pi}\right)^2 \frac{G_F^2 m_b^5 |V_{ts}^* V_{tb}|^2}{48\pi^3} (1-\hat{s})^2 \times \left[(1+2\hat{s}) (|C_9^{eff}|^2 + |C_{10}^{eff}|^2) + 4 \left(1 + \frac{2}{\hat{s}}\right) |C_7^{eff}|^2 + 12 \text{Re}(C_7^{eff} C_9^{eff*}) \right]$$

Particularly sensitive: backward-forward asymmetry in $K^* l^+ l^-$


January 5, 2006

University of Dortmund

Peter Krizan, Ljubljana



$$A_{FB}(q^2) = \frac{\int_0^1 \frac{d^2\Gamma}{dq^2 d\cos\theta} d\cos\theta - \int_{-1}^0 \frac{d^2\Gamma}{dq^2 d\cos\theta} d\cos\theta}{\int_0^1 \frac{d^2\Gamma}{dq^2 d\cos\theta} d\cos\theta + \int_{-1}^0 \frac{d^2\Gamma}{dq^2 d\cos\theta} d\cos\theta}$$




[γ^* and Z^* contributions in $B \rightarrow K^* l l$ interfere and give rise to forward-backward asymmetries c.f. $e^+e^- \rightarrow \mu^+ \mu^-$]

Unbinned fit to the variables q^2 (di-lepton invariant mass) and $\cos(\theta)$ for $B \rightarrow K^* l l$ data




2005: Sample used for $A_{FB}(B \rightarrow K^* l l)(q^2)$



Sample for $B \rightarrow K^* l l$ events 113 ± 13
 $B \rightarrow K l l$ control sample 96 ± 12
 Consistent with flat

$$P(q^2, \cos\theta; A_0/A_T, A_{10}/A_T) = f_{sig} \epsilon_{sig}(q^2, \cos\theta) \frac{d^2\Gamma}{dq^2 d\cos\theta}(q^2, \cos\theta) / N_{sig} + f_{ctct} \epsilon_{ctct}(q^2, \cos\theta) \frac{d^2\Gamma}{dq^2 d\cos\theta}(q^2, \cos\theta) / N_{ctct} + f_{fctct} \epsilon_{fctct}(q^2, \cos\theta) \frac{d^2\Gamma}{dq^2 d\cos\theta}(q^2, -\cos\theta) / N_{fctct} + f_{X,tl} P_{X,tl}(q^2, \cos\theta) + f_{all} \{ (1 - f_{K^*lh}) P_{all}(q^2, \cos\theta) + f_{K^*lh} P_{K^*lh}(q^2, \cos\theta) \} + f_{K^*hh} P_{K^*hh}(q^2, \cos\theta) + f_{\psi} P_{\psi}(q^2, \cos\theta)$$


Treat $q^2, \cos(\theta)$ dependence of bkg.





Constraints on Wilson coefficients from $A_{FB}(B \rightarrow K^* l l)(q^2)$

Projections of the full fit to $q^2, \cos(\theta)$



Integrated FB asymmetry

$$A_{FB}(B \rightarrow K^* l^- l^+) = 0.50 \pm 0.12 \pm 0.02; (3.4\sigma)$$

control sample:

$$A_{FB}(B \rightarrow K^+ l^- l^+) = 0.10 \pm 0.14 \pm 0.01$$

Observed integrated A_{FB} rules out some radical New Physics Models with incorrect signs/magnitudes of C_9 and C_{10}



January 5, 2006

University of Dortmund

Peter Krizan, Ljubljana



Results of the unbinned fit to q^2 and $\cos(\theta)$ distributions for ratios of Wilson coefficients.



$|A_7|$ constrained from $b \rightarrow s \gamma$ to be close to SM

A_9/A_7

	negative A_7	positive A_7
A_9/A_7	$-15.3^{+3.4}_{-4.8} \pm 1.1$	$-16.3^{+3.7}_{-5.7} \pm 1.4$
A_{10}/A_7	$10.3^{+5.2}_{-3.5} \pm 1.8$	$11.1^{+6.0}_{-3.9} \pm 2.4$

Ref: hep-ex/0508009


$$-1401 < A_9 A_{10}/A_7^2 < -26.4 \text{ at } 95\% \text{ C.L.}$$

Peter Krizan, Ljubljana




$A_{FB}(B \rightarrow K^* l^+ l^-)[q^2]$ at Super B Factory

- Assume 1 year of running at 5×10^{35} nb/sec
- \rightarrow 5/ab integrated luminosity, 10 billion B mesons
- $\Delta A_9/A_9 \sim 11\%$, $\Delta A_{10}/A_{10} \sim 13\%$
- A_7 fixed to SM value



Determine location of the zero crossing precisely with 50 ab^{-1}



January 5, 2006

University of Dortmund

Peter Krizan, Ljubljana



Fundamental Questions in Flavor Physics

Are there New Physics Phases and New sources of CP Violation Beyond the SM ?

Experiments: $b \rightarrow s$ CPV, compare CPV angles from tree and loops

Are there new operators with quarks enhanced by New Physics ?

Experiments: $A_{FB}(B \rightarrow K^* l^+ l^-)$, $B \rightarrow K \pi$ rates and asymmetries

Are there right-handed currents ?

Experiments: $b \rightarrow s \gamma$ CPV, $B \rightarrow V V$ triple-product asymmetries

Are there new flavor changing neutral currents ?

Experiments: $b \rightarrow s \nu \nu$, D - \bar{D} mixing+CPV+rare, $\tau \rightarrow \mu \gamma$

These questions can only be answered at a Super B Factory.

January 5, 2006

University of Dortmund

Peter Krizan, Ljubljana




Super B Factory Motivation

- Physics beyond the Standard Model (SM) must exist.
 - finite m_ν
 - gravity
- If the LHC finds New Physics at the TeV scale,
 - its flavor structure must be examined experimentally. A super B factory is the best tool for this purpose.



cf. Physics of top quark
 First estimate of mass: BB mixing → ARGUS
 Direct production, Mass, width etc. → CDF/D0
 Off-diagonal couplings, phase → BaBar/Belle



- If the LHC finds nothing but a SM-like Higgs,
 - searching for deviations from the SM in flavor physics will be one of the best ways to find new physics.

January 5, 2006

University of Dortmund


Peter Krizan, Ljubljana



New physics effects in B physics


Different New Physics scenarios and their effects in B decays.

G.Hiller





A Broad Unbiased Approach to New Physics




January 5, 2006

University of Dortmund

Peter Krizan, Ljubljana



CPV in $b \rightarrow s$ and diagnosis of new physics



Expected precision at $5ab^{-1}$

Many other examples of using correlations to distinguish new physics scenarios have been examined.

T.Goto, Y.Okada, Y.Shimizu, T.Shindou, M.Tanaka (2002, 2004) + SuperKEKB LoI

January 5, 2006


University of Dortmund

Peter Krizan, Ljubljana



Super-B and LHCb: complementary

- **Clean environment** → measurements that no other experiment can perform. Examples: CPV in $B \rightarrow \phi K^0$, $B \rightarrow \eta' K^0$ for new phases, $B \rightarrow K_S \pi^0 \gamma$ for right-handed currents.
- **"B-meson beam" technique** → access to new decay modes. Example: discover $B \rightarrow K \nu \nu$.
- **Measure new types of asymmetries**
Example: forward-backward asymmetry in $b \rightarrow s \mu \mu$, *see*
- **Rich, broad physics program including B, τ and charm physics**
Examples: searches for $\tau \rightarrow \mu \gamma$ and D - D mixing with unprecedented sensitivity.



January 5, 2006



University of Dortmund



Super B Factory at KEK

New Beam pipe

Ante-chamber & solenoid coils to reduce photo-electron clouds



More RF power

Damping ring

Linac upgrade $L = 4 \times 10^{35} / \text{cm}^2 / \text{s}$

January 5, 2006


University of Dortmund

Peter Krizan, Ljubljana




Crab crossing in the near future

- Crab crossing may increase the beam-beam parameter up to 0.19 !



K. Ohmi

- Superconducting crab cavities are now being tested, will be installed in KEKB around **March 2006**.





Requirements for the Super B detector

Critical issues at $L = 4 \times 10^{35}/\text{cm}^2/\text{sec}$

- ▶ **Higher background ($\times 20$)**
 - radiation damage and occupancy
 - fake hits and pile-up noise in the EM
- ▶ **Higher event rate ($\times 10$)**
 - higher rate trigger, DAQ and computing
- ▶ **Require special features**
 - low $p \mu$ identification $\leftarrow s \mu \mu$ recon. eff.
 - hermeticity $\leftarrow \nu$ "reconstruction"

Possible solution:

- ▶ Replace inner layers of the vertex detector with a silicon strip detector.
- ▶ Replace inner part of the central tracker with a silicon strip detector.
- ▶ Better particle identification device
- ▶ Replace endcap calorimeter by pure CsI.
- ▶ Faster readout electronics and computing system.




January 5, 2006

University of Dortmund


Peter Krizan, Ljubljana



Belle Upgrade for Super-B



PID upgrade in the endcap



improve K/π separation in the forward (high mom.) region for few-body decays of B's

good K/π separation for $b \rightarrow d\gamma$, $b \rightarrow s\gamma$

improve purity in fully reconstructed B decays

low momentum ($< 1\text{GeV}/c$) $e/\mu/\pi$ separation ($B \rightarrow K\ell\ell$)

keep high the efficiency for tagging kaons


January 5, 2006

University of Dortmund

Peter Krizan, Ljubljana



Proximity focusing RICH in the forward region



K/ π separation at 4 GeV/c
 $\theta_c(\pi) \sim 308$ mrad ($n = 1.05$)
 $\theta_c(\pi) - \theta_c(K) \sim 23$ mrad

$d\theta_c(\text{meas.}) = \sigma_0 \sim 13$ mrad
With 20mm thick aerogel and
6mm PMT pad size

$\rightarrow 6\sigma$ separation with $N_{pe} \sim 10$

January 5, 2006

University of Dortmund


Peter Krizan, Ljubljana



Beam test: Cherenkov angle resolution and number of photons

Beam test results with 2cm thick aerogel tiles:

$>4\sigma$ K/ π separation



\rightarrow Number of photons has to be increased.

January 5, 2006

University of Dortmund




Peter Krizan, Ljubljana



How to increase the number of photons?

What is the optimal radiator thickness?

Use beam test data on σ_0 and N_{pe}



Minimize the error per track:

$$\sigma_{track} = \sigma_0 / \sqrt{N_{pe}}$$

Optimum is close to 2 cm

January 5, 2006


University of Dortmund



Radiator with multiple refractive indices

How to increase the number of photons without degrading the resolution?

-> stack two tiles with different refractive indices: "focusing" configuration



NIM A548 (2005) 383

January 5, 2006




University of Dortmund

Peter Krizan, Ljubljana






Focusing configuration – data

4cm aerogel single index



2+2cm aerogel




January 5, 2006



Tiling of the radiator

Minimize photon yield losses at the aerogel tile boundary: hexagonal tiling scheme



- Cut into hexagonal shape from a square block
- Machining device: use "water-jet" thanks to the hydrophobic nature


January 5, 2006

University of Dortmund

Peter Krizan, Ljubljana



KEKB Collider Upgrade Scenario




L_{peak} ($\text{cm}^{-2}\text{s}^{-1}$)	1.6×10^{34}	\rightarrow	5×10^{34}	\rightarrow	5×10^{35}
L_{int}	469fb^{-1}	\rightarrow	$\sim 1 \text{ab}^{-1}$	\rightarrow	$\sim 10 \text{ab}^{-1}$
	January 5, 2006		University of Dortmund		Peter Krizan, Ljubljana



Yamauchi's Schedule for Super B


A Super B proposal was submitted to MEXT in August 2005.
 KEBK/Belle project receives a grade of S(i.e. A+) in gov. review
 A search for a new KEK laboratory director is underway.



January 5, 2006 University of Dortmund Peter Krizan, Ljubljana



Yamauchi's budget for Super B



January 5, 2006

University of Dortmund

Peter Krizan, Ljubljana



Summary 1

- Observation of direct CP violation in $B^0 \rightarrow \pi^+\pi^-$ and $K^+\pi^-$ decays, evidence in $B^- \rightarrow \rho^0 K^-$
- CP violation in $b \rightarrow s$ transitions remains below SM expectation, but **statistically limited**.
- Forward-backward asymmetry (A_{FB}) in $b \rightarrow s l^+ l^-$ is becoming another powerful tool to search for physics beyond SM.
- We are entering an exciting phase of **precision measurements**.

.... and there are much more interesting results, but could not be covered in this talk!

January 5, 2006

University of Dortmund

Peter Krizan, Ljubljana



Summary 2

- B factories have proven to be an excellent tool for flavour physics
- Reliable long term operation, constant improvement of the performance.
- Short term plan: increase luminosity **x3** by a crab cavity
- Major upgrade in 2009-10 -> Super B factory, **L x30**
- Essentially a new project, all components have to be replaced, plans exist (LoI), nothing is frozen...
- Expect a new, exciting era of discoveries, complementary to LHC
- Do not miss the chance to be part of it..

January 5, 2006

University of Dortmund

Peter Krizan, Ljubljana