



# HERA-B RICH: performance and physics impact

**Marko Starič**


Institut Jožef Stefan, Ljubljana, Slovenia

**Playa del Carmen, 30 Nov. - 5 Dec. 2004**  
**5<sup>th</sup> International Workshop on RICH Counters**

1. Introduction
2. RICH performance
3. RICH in physics analysis
4. Summary



## The HERA-B experiment

- ◆ Fixed target experiment at HERA proton beam (920 GeV/c)
  - ▷ High rate forward spectrometer (<40 MHz interaction rate)
  - ▷ Wire targets of different material in the beam halo
  - ▷ High resolution vertexing
  - ▷ Very good particle ID for  $e$ ,  $\mu$ ,  $\pi$ ,  $K$  and  $p$
  - ▷ Hardware track trigger for lepton pairs ( $J/\psi \rightarrow l^+l^-$ )




## The HERA-B RICH

- ◆  $108m^3$  of  $C_4F_{10}$  ( $\gamma_t = 19.1$ )
- ◆ Two spherical mirrors, tilted by  $9^0$ ,  $f = 5.7m$
- ◆ Photon detector with Hamamatsu multi-anode PMT's



## Parameters of the RICH

- ◆ Čerenkov angle for  $\beta = 1$  particle: 52 mrad
- ◆ Number of photons per  $\beta = 1$  particle: 32
- ◆ Figure of merit  $N_0$ :  $42\text{cm}^{-1}$
- ◆ Single photon angular resolution:
  - ▷ 16 channel PMT region:  $(0.7 \oplus 3.5/p)$  mrad
  - ▷ 4 channel PMT region:  $(1.0 \oplus 3.5/p)$  mrad
  - ▷ including track error: 1.2 mrad (mean), 0.8 mrad (above 40GeV/c)



$\pi - K$  separation:  
5 - 50 GeV/c

$\pi - p$  separation:  
5 - 100 GeV/c

$K - p$  separation:  
10 - 100 GeV/c

## Identification efficiencies

- ◆ Extended maximum likelihood method used (reported on RICH2002, Pylos)  
→ normalized likelihood probabilities for  $e, \mu, \pi, K, p$
- ◆ Particle selection made by applying a cut on the appropriate likelihood
- ◆ Identification efficiencies measured on real data by using the decays:


$$K_s^0 \rightarrow \pi^+ \pi^-$$

$$\phi \rightarrow K^+ K^-$$


$$\Lambda(\bar{\Lambda}) \rightarrow p\pi^-(\bar{p}\pi^+)$$

### Pion identification


$$lre + lr\mu + lr\pi > 0.05$$



Pions



Kaons




Protons



## Kaon identification

$lrk > 0.50$




Pions

Kaons

Protons

$lrk > 0.30$





Pions

Kaons


Protons

## Proton identification

 $lrp > 0.45$ *Pions**Kaons**Protons* $lrp > 0.90$ *Pions**Kaons**Protons*

## RICH in Physics Analysis

- ◆ RICH is playing a crucial role in physics analysis, where decay products include kaons and/or protons.
- ◆ At hadron collisions:
  - large particle multiplicities (10 - 30 charged tracks)
  - mostly pions (70% - 80%)
    - large combinatorial background
- ◆ Good particle identification:
  - kills combinatorial background
  - removes reflections





## Another Two Examples





$\Lambda(1520) \rightarrow pK^-$

short lived  
(decays inside target)



$\Omega^- \rightarrow \Lambda K^-$

long lived  
(decays far away)






## Physics at HERA-B




- ◆ Mainly devoted to studies of particle production in proton-nucleus collisions
- ◆ Two large data samples were collected during 2002/2003 run period
  - di-lepton trigger data: 150M events, 300k  $J/\psi$
  - minimum bias data (interaction trigger): 210M events
- ◆ Three analysis will be briefly presented:
  - production of vector mesons  $\phi$  and  $K^{*0}$
  - production of open charm
  - search for  $\Theta^+$  and  $\Xi^{--}$  pentaquarks
- ◆ Minimum bias data used (C, Ti and W target)

## $K^{*0}(892)$ and $\phi(1020)$ production


- ◆ Minimum bias data (150M events)
- ◆ Three target materials: C, Ti, W
- ◆ Data selection:
  - vertex reconstruction
  - kaon identification



$$N_{K^{*0}} = 540k$$



$$N_{\bar{K}^{*0}} = 420k$$





$$N_\phi = 51k$$

# $K^{*0}(892)$ and $\phi(1020)$ production



Preliminary results for:


- ◆ total production cross sections  $\sigma_{pC}$ ,  $\sigma_{pTi}$  and  $\sigma_{pW}$
- ◆ single differential production cross sections  $d\sigma/dy$  and  $d\sigma/dp_T^2$
- ◆ A-dependence of cross sections:  $\sigma_{pA} = \sigma_{pN} A^\alpha$ 
  - ▷ proton-nucleon cross section  $\sigma_{pN}$  and exponent  $\alpha$
- ◆ Cronin effect observed for the first time in the production of  $K^{*0}$  and  $\phi$  (exponent  $\alpha$  increases with  $p_T$ )



## Open Charm


- ◆ Minimum bias data (200M events)
- ◆ Charm cross section 3 orders of magnitude smaller than inelastic
- ◆ Large boost ( $\gamma = 22$ ): D mesons decay several mm from target
- ◆ Data selection:
  - vertex reconstruction (resolution  $\sigma_{\Delta z} \approx 500\mu\text{m}$ )
  - detached vertex requirement
  - kaon and pion identification

$D^0 \rightarrow K^-\pi^+$




$$N_{D^0} = 194 \pm 20$$

$D^+ \rightarrow K^-\pi^+\pi^+$



$$N_{D^+} = 92 \pm 11$$

$D^{*+} \rightarrow D^0\pi^+$





$$N_{D^{*+}} = 49 \pm 10$$

## Open charm

Preliminary results for:

- ◆ total production cross sections  $\sigma_{pC}$ ,  $\sigma_{pTi}$ ,  $\sigma_{pW}$  and  $\sigma_{pN}$
- ◆ D meson production ratios:  $D^+/D^0$  and  $D^{*+}/D^0$




## Search for pentaquarks

Motivation:

- ◆ Recently more than 10 experiments have reported the observation of narrow state in either  $nK^+$  or  $pK_S^0$  decay channels at  $1540 \text{ MeV}/c^2$ .
- ◆ Another candidate resonance have been found in  $\Xi^-\pi^-$  and  $\Xi^-\pi^+$  decay channel at  $1862 \text{ MeV}/c^2$ .
- ◆ HERA-B collected large minimum bias data sample
- ◆ Statistics of the relevant reconstructed signals

signal	statistics	$\sigma (\text{MeV}/c^2)$
$K_S^0$	4.9M	4.9
$\Lambda$ [c.c]	1.1M [520k]	1.6
$\Lambda(1520)$ [c.c]	3.5k [2.1k]	2.3
$\Xi^-$ [c.c]	12k [8.2k]	2.6
$\Xi(1530)^0$ [c.c]	1.4k [940]	2.9



# Search for pentaquarks





## Particle ID:

- ◆  $\Theta^+ \rightarrow p K_S^0$  strong cut on proton likelihood (>0.95)
  - ▷ mis-identification <1%
- ◆  $\Xi^{--} \rightarrow \Xi^- \pi^-$  week cut on pion likelihood (>0.05)
  - ▷ reduces background from K and p

## Results:

- ◆ No evidence for PQ narrow signals found
  - ▷ Upper limits for production cross sections
  - ▷ UL for yield ratio to  $\Lambda(1520)$  and  $\Xi(1530)^0$
- ◆ Published in *Phys.Rev.Lett.93, 212003(2004)*



## Summary

- ◆ Performance of the HERA-B RICH and its impact on physics analysis have been presented.
- ◆ With the RICH we can identify pions, kaons and protons essentially in the entire kinematic range of our experiment, with the efficiency as large as 90% and with the mis-identification probability at the 1% level.
- ◆ By kaon and proton identification the combinatorial background is in some cases reduced by more than 3 orders of magnitude.
- ◆ Three physics analysis were shown, which wouldn't be possible without high performance of the RICH particle identification system.
- ◆ The HERA-B experiment finished data taking in spring 2003. The spectrometer (and RICH) has been dismounted.

