



CP violation

Part 2: CP violation primer

Peter Križan

University of Ljubljana and J. Stefan Institute

May 17-25, 2005

Course at University of Barcelona

Peter Križan, Ljubljana



Contents

CP violation in the B system
Standard Model predictions
CP violation in the K system

May 17-25, 2005

Course at University of Barcelona

Peter Križan, Ljubljana



Time evolution in the B system

An arbitrary linear combination of the neutral B-meson flavor eigenstates

$$a|B^0\rangle + b|\bar{B}^0\rangle$$

is governed by a time-dependent Schrodinger equation

$$i \frac{d}{dt} \begin{pmatrix} a \\ b \end{pmatrix} = H \begin{pmatrix} a \\ b \end{pmatrix} = (M - i\Gamma) \begin{pmatrix} a \\ b \end{pmatrix}$$

M and Γ are 2x2 Hermitian matrices. *CPT* invariance $\rightarrow H_{11} = H_{22}$

May 17-25, 2005

Course at University of Barcelona

Peter Krizan, Ljubljana



Time evolution in the B system

The light B_L and heavy B_H mass eigenstates are given by

$$|B_L\rangle = p|B^0\rangle + q|\bar{B}^0\rangle$$

$$|B_H\rangle = p|B^0\rangle - q|\bar{B}^0\rangle$$

With the eigenvalue differences

$$\Delta m_B = m_H - m_L, \Delta\Gamma_B = \Gamma_H - \Gamma_L$$

Which are related to the M and Γ matrix elements

$$(\Delta m_B)^2 - \frac{1}{4}(\Delta\Gamma_B)^2 = 4(|M_{12}|^2 - \frac{1}{4}|\Gamma_{12}|^2)$$

$$\Delta m_B \Delta\Gamma_B = 4 \operatorname{Re}(M_{12} \Gamma_{12}^*)$$

May 17-25, 2005

Course at University of Barcelona

Peter Krizan, Ljubljana



The ratio p/q is

$$\frac{q}{p} = -\frac{\Delta m_B - \frac{i}{2} \Delta \Gamma_B}{2(M_{12} - \frac{i}{2} \Gamma_{12})} = -\frac{2(M_{12}^* - \frac{i}{2} \Gamma_{12}^*)}{\Delta m_B - \frac{i}{2} \Delta \Gamma_B}$$

What do we know about Δm_B and $\Delta \Gamma_B$?

$x_d = \Delta m_B / \Gamma_B = 0.73 \pm 0.05$ well measured

$\Delta \Gamma_B / \Gamma_B$ not measured, expected $O(0.01)$, due to decays common to B and anti-B - $O(0.001)$.

$\rightarrow \Delta \Gamma_B \ll \Delta m_B$

May 17-25, 2005

Course at University of Barcelona

Peter Krizan, Ljubljana



Since $\Delta \Gamma_B \ll \Delta m_B$

$$\Delta m_B = 2|M_{12}|$$

$$\Delta \Gamma_B = 2 \operatorname{Re}(M_{12} \Gamma_{12}^*) / |M_{12}|$$

and

$$\frac{q}{p} = -\frac{|M_{12}|}{M_{12}}$$

or to next order

$$\frac{q}{p} = -\frac{|M_{12}|}{M_{12}} \left[1 - \frac{1}{2} \operatorname{Im} \left(\frac{\Gamma_{12}}{M_{12}} \right) \right]$$

May 17-25, 2005

Course at University of Barcelona

Peter Krizan, Ljubljana



Any B state can then be written as an admixture of the states B_H and B_L , and the amplitudes of this admixture evolve in time

$$a_H(t) = a_H(0)e^{-iM_H t} e^{-\Gamma_H t/2}$$


$$a_L(t) = a_L(0)e^{-iM_L t} e^{-\Gamma_L t/2}$$

A B^0 state created at $t=0$ (denoted by B^0_{phys}) has $a_H(0) = a_L(0) = 1/(2p)$; an anti-B at $t=0$ ($\text{anti-}B^0_{\text{phys}}$) has $a_H(0) = a_L(0) = 1/(2q)$

May 17-25, 2005

Course at University of Barcelona

Peter Krizan, Ljubljana



B^0 at $t=0$, evolution in time

Full line: B^0 , dotted: \bar{B}^0

T: in units of $\tau=1/\Gamma$

May 17-25, 2005

Course at University of Barcelona

Peter Krizan, Ljubljana



Time evolution:

$$|B_{phys}^0(t)\rangle = g_+(t)|B^0\rangle + (q/p)g_-(t)|\bar{B}^0\rangle$$

$$|\bar{B}_{phys}^0(t)\rangle = (p/q)g_-(t)|B^0\rangle + g_+(t)|\bar{B}^0\rangle$$

with

$$g_+(t) = e^{-iMt} e^{-\Gamma t/2} \cos(\Delta m t / 2)$$

$$g_-(t) = e^{-iMt} e^{-\Gamma t/2} i \sin(\Delta m t / 2)$$

$$M = (M_H + M_L)/2$$

May 17-25, 2005

Course at University of Barcelona

Peter Krizan, Ljubljana



CP violation: three types

Define decay amplitudes of B and anti-B to the same final state f

$$A_f = \langle f | H | B^0 \rangle$$

$$\bar{A}_f = \langle f | H | \bar{B}^0 \rangle$$

Define also parameter λ

$$\lambda = \frac{q}{p} \frac{\bar{A}_f}{A_f}$$

Three types of CP violation (CPV):

$$\left. \begin{array}{l} \text{CP in decay: } |\bar{A}/A| \neq 1 \\ \text{CP in mixing: } |q/p| \neq 1 \end{array} \right\} |\lambda| \neq 1$$

CP in interference between mixing and decay: even if $|\lambda| = 1$ if only $\text{Im}(\lambda) \neq 0$

May 17-25, 2005

Course at University of Barcelona

Peter Krizan, Ljubljana



CP violation in decay

\mathcal{CP} in decay: $|\bar{A}/A| \neq 1$
(and of course also $|\lambda| \neq 1$)

$$a_f = \frac{\Gamma(B^+ \rightarrow f, t) - \Gamma(B^- \rightarrow \bar{f}, t)}{\Gamma(B^+ \rightarrow f, t) + \Gamma(B^- \rightarrow \bar{f}, t)} = \frac{1 - |\bar{A}/A|^2}{1 + |\bar{A}/A|^2}$$

Also possible for neutral B.

May 17-25, 2005

Course at University of Barcelona

Peter Krizan, Ljubljana



CP violation in decay

CPV in decay: $|\bar{A}/A| \neq 1$: how do we get there?

In general, A is a sum of amplitudes with strong phases δ_i and weak phases ϕ_i . The amplitudes for anti-particles have same strong phases and opposite weak phases ->

$$A_f = \sum_i A_i e^{i(\delta_i + \phi_i)}$$

$$\bar{A}_f = \sum_i A_i e^{i(\delta_i - \phi_i)}$$

$$\left| \frac{\bar{A}_f}{A_f} \right| = \left| \frac{\sum_i A_i e^{i(\delta_i - \phi_i)}}{\sum_i A_i e^{i(\delta_i + \phi_i)}} \right|$$

$$|A_f|^2 - |\bar{A}_f|^2 = \sum_{i,j} A_i A_j \sin(\phi_i - \phi_j) \sin(\delta_i - \delta_j)$$

CPV in decay: need at least two interfering amplitudes with different weak and strong phases.

May 17-25, 2005

Course at University of Barcelona

Peter Krizan, Ljubljana



CP violation in mixing

CP in mixing: $|q/p| \neq 1$

(again $|\lambda| \neq 1$)

In general: probability for a B to turn into an anti-B can be different from the probability for an anti-B to turn into a B.

$$\begin{aligned}
 |B_{phys}^0(t)\rangle &= g_+(t)|B^0\rangle + (q/p)g_-(t)|\bar{B}^0\rangle \\
 |\bar{B}_{phys}^0(t)\rangle &= (p/q)g_-(t)|B^0\rangle + g_+(t)|\bar{B}^0\rangle
 \end{aligned}$$

Example: semileptonic decays:

$$\begin{aligned}
 \langle l^- \nu X | H | B_{phys}^0(t) \rangle &= (q/p)g_-(t)A^* \\
 \langle l^+ \nu X | H | \bar{B}_{phys}^0(t) \rangle &= (p/q)g_-(t)A
 \end{aligned}$$

May 17-25, 2005

Course at University of Barcelona

Peter Kriz̃an, Ljubljana



CP violation in mixing

$$\begin{aligned}
 a_{sl} &= \frac{\Gamma(\bar{B}_{phys}^0(t) \rightarrow l^+ \nu X) - \Gamma(B_{phys}^0(t) \rightarrow l^- \nu X)}{\Gamma(\bar{B}_{phys}^0(t) \rightarrow l^+ \nu X) + \Gamma(B_{phys}^0(t) \rightarrow l^- \nu X)} = \\
 &= \frac{1 - |q/p|^4}{1 + |q/p|^4}
 \end{aligned}$$

-> Small, since to first order $|q/p| \sim 1$. Next order:

$$\frac{q}{p} = -\frac{|M_{12}|}{M_{12}} \left[1 - \frac{1}{2} \text{Im} \left(\frac{\Gamma_{12}}{M_{12}} \right) \right]$$

Expect O(0.01) effect in semileptonic decays

May 17-25, 2005

Course at University of Barcelona


Peter Kriz̃an, Ljubljana



CP violation in the interference between decays with and without mixing

CP violation in the interference between mixing and decay to a state accessible in both B^0 and anti- B^0 decays

For example: a CP eigenstate f_{CP} like $\pi^+ \pi^-$



$$\lambda = \frac{q}{p} \frac{\bar{A}_f}{A_f}$$

We can get CP violation if $\text{Im}(\lambda) \neq 0$, even if $|\lambda| = 1$

May 17-25, 2005

Course at University of Barcelona

Peter Kriz̃an, Ljubljana



CP violation in the interference between decays with and without mixing

Decay rate asymmetry:

$$a_{f_{CP}} = \frac{P(\bar{B}^0 \rightarrow f_{CP}, t) - P(B^0 \rightarrow f_{CP}, t)}{P(\bar{B}^0 \rightarrow f_{CP}, t) + P(B^0 \rightarrow f_{CP}, t)}$$

Decay rate: $P(B^0 \rightarrow f_{CP}, t) \propto \left| \langle f_{CP} | H | B_{phys}^0(t) \rangle \right|^2$

Decay amplitudes vs time:

$$\begin{aligned} \langle f_{CP} | H | B_{phys}^0(t) \rangle &= g_+(t) \langle f_{CP} | H | B^0 \rangle + (q/p) g_-(t) \langle f_{CP} | H | \bar{B}^0 \rangle \\ &= g_+(t) A_{f_{CP}} + (q/p) g_-(t) \bar{A}_{f_{CP}} \end{aligned}$$

$$\begin{aligned} \langle f_{CP} | H | \bar{B}_{phys}^0(t) \rangle &= (p/q) g_-(t) \langle f_{CP} | H | B^0 \rangle + g_+(t) \langle f_{CP} | H | \bar{B}^0 \rangle \\ &= (p/q) g_-(t) A_{f_{CP}} + g_+(t) \bar{A}_{f_{CP}} \end{aligned}$$

May 17-25, 2005

Course at University of Barcelona

Peter Kriz̃an, Ljubljana



CP violation in the interference between decays with and without mixing

$$a_{f_{CP}} = \frac{P(\bar{B}^0 \rightarrow f_{CP}, t) - P(B^0 \rightarrow f_{CP}, t)}{P(\bar{B}^0 \rightarrow f_{CP}, t) + P(B^0 \rightarrow f_{CP}, t)} = \lambda = \frac{q}{p} \frac{\bar{A}_f}{A_f}$$

$$= \frac{(1 - |\lambda_{f_{CP}}|^2) \cos(\Delta mt) - 2 \operatorname{Im}(\lambda_{f_{CP}}) \sin(\Delta mt)}{1 + |\lambda_{f_{CP}}|^2}$$

Non-zero effect if $\operatorname{Im}(\lambda) \neq 0$,
even if $|\lambda| = 1$

If in addition $|\lambda| = 1 \rightarrow$

$$a_{f_{CP}} = -\operatorname{Im}(\lambda_{f_{CP}}) \sin(\Delta mt)$$

May 17-25, 2005

Course at University of Barcelona

Peter Krizan, Ljubljana



CP violation in the interference between decays with and without mixing

One more form for λ :

$$\lambda_{f_{CP}} = \frac{q}{p} \frac{\bar{A}_{f_{CP}}}{A_{f_{CP}}} = \eta_{f_{CP}} \frac{q}{p} \frac{\bar{A}_{f_{CP}}}{A_{f_{CP}}}$$

$\eta_{f_{CP}} = \pm 1$ CP parity of f_{CP}

\rightarrow we get one more (-1) sign when comparing
asymmetries in two states with opposite CP parity

$$a_{f_{CP}} = -\operatorname{Im}(\lambda_{f_{CP}}) \sin(\Delta mt)$$

May 17-25, 2005

Course at University of Barcelona

Peter Krizan, Ljubljana



B and anti-B from the Y(4s)

B and anti-B from the Y(4s) decay are in a $l=1$ state.

They cannot mix independently (either BB or anti-B anti-B states are forbidden with $l=1$ due to Bose symmetry).

After one of them decays, the other evolves independently ->

-> only time differences between one and the other decay matter (for mixing).

Assume

- one decays to a CP eigenstate f_{CP} (e.g. $\pi\pi$ or $J/\psi K_S$) at time $t_{f_{CP}}$ and
 - the other at t_{tag} to a flavor-specific state f_{tag} (=state only accessible to a B^0 and not to a anti- B^0 (or vice versa), e.g. $B^0 \rightarrow D^0\pi$, $D^0 \rightarrow K^-\pi^+$)
- also known as 'tag' because it tags the flavour of the B meson it comes from

May 17-25, 2005

Course at University of Barcelona

Peter Krizan, Ljubljana



Time evolution for B and anti-B from the Y(4s)

The time evolution for the B anti-B pair from Y(4s) decay

$$R(t_{tag}, t_{f_{CP}}) = e^{-\Gamma(t_{tag} + t_{f_{CP}})} \left| \overline{A_{tag}} \right|^2 \left| A_{f_{CP}} \right|^2$$

$$\left[1 + \left| \lambda_{f_{CP}} \right|^2 + \cos[\Delta m(t_{tag} - t_{f_{CP}})] (1 - \left| \lambda_{f_{CP}} \right|^2) \right.$$

$$\left. - 2 \sin(\Delta m(t_{tag} - t_{f_{CP}})) \text{Im}(\lambda_{f_{CP}}) \right]$$

with $\lambda_{f_{CP}} = \frac{q}{p} \frac{\overline{A}_{f_{CP}}}{A_{f_{CP}}}$

-> in asymmetry measurements at Y(4s) we have to use $t_{tag} - t_{f_{CP}}$ instead of absolute time t .

May 17-25, 2005


Course at University of Barcelona

Peter Krizan, Ljubljana




Decay rate to f_{CP}

Incoherent production
(e.g. hadron collider)



coherent production
at $Y(4s)$



May 17-25, 2005


Course at University of Barcelona

Peter Krizan, Ljubljana



CP violation in SM

CP violation: consequence of the
Cabibbo-Kobayashi-Maskawa (CKM)
quark mixing matrix



$$V_{CKM} = \begin{pmatrix} V_{ud} & V_{us} & V_{ub} \\ V_{cd} & V_{cs} & V_{cb} \\ V_{td} & V_{ts} & V_{tb} \end{pmatrix}$$

May 17-25, 2005

Course at University of Barcelona

Peter Krizan, Ljubljana



CP violation in SM

$$\mathcal{L} = V_{ij} \bar{U}_i \gamma^\mu (1 - \gamma_5) D_j W_\mu^+ + V_{ij}^* \bar{D}_i \gamma^\mu (1 - \gamma_5) U_j W_\mu^-$$

$\Downarrow CP$

$$\mathcal{L}_{CP} = V_{ij} \bar{D}_i \gamma^\mu (1 - \gamma_5) U_j W_\mu^- + V_{ij}^* \bar{U}_i \gamma^\mu (1 - \gamma_5) D_j W_\mu^+$$

If $V_{ij} = V_{ij}^* \quad \blacktriangleright \quad \mathcal{L} = \mathcal{L}_{CP} \quad \blacktriangleright \quad CP \text{ is conserved}$

May 17-25, 2005

Course at University of Barcelona

Peter Krizán, Ljubljana



CKM matrix

3x3 orthogonal matrix: 3 parameters - angles

3x3 unitary matrix: 18 parameters, 9 conditions = 9 free parameters, 3 angles and 6 phases

6 quarks: 5 relative phases can be transformed away (by redefining the quark fields)

1 phase left -> the matrix is in general complex

$$V_{CKM} = \begin{pmatrix} c_{12}c_{13} & s_{12}c_{13} & s_{13}e^{-i\delta} \\ -s_{12}c_{13} - c_{12}s_{23}s_{13}e^{i\delta} & c_{12}c_{23} - s_{12}s_{23}s_{13}e^{i\delta} & s_{23}c_{13} \\ s_{12}s_{23} - c_{12}c_{23}s_{13}e^{i\delta} & -c_{12}s_{23} - s_{12}c_{23}s_{13}e^{i\delta} & c_{23}c_{13} \end{pmatrix}$$

$s_{12} = \sin\theta_{12}$, $c_{12} = \cos\theta_{12}$ etc.


May 17-25, 2005

Course at University of Barcelona

Peter Krizán, Ljubljana




CKM matrix



Transitions between members of the same family more probable (=thicker lines) than others

-> CKM: almost a diagonal matrix, but not completely ->



May 17-25, 2005

Course at University of Barcelona

Peter Kriz̃an, Ljubljana



CKM matrix

Almost a diagonal matrix, but not completely ->

Wolfenstein parametrisation: expand in the parameter λ ($=\sin\theta_c=0.22$)

A , ρ and η : all of order one

$$V = \begin{pmatrix} 1 - \frac{\lambda^2}{2} & \lambda & A\lambda^3(\rho - i\eta) \\ -\lambda & 1 - \frac{\lambda^2}{2} & A\lambda^2 \\ A\lambda^3(1 - \rho - i\eta) & -A\lambda^2 & 1 \end{pmatrix} + O(\lambda^4)$$

May 17-25, 2005

Course at University of Barcelona

Peter Kriz̃an, Ljubljana



CKM matrix

define $s_{12} \equiv \lambda, s_{23} \equiv A\lambda^2, s_{13}e^{-i\delta} \equiv A\lambda^3(\rho - i\eta)$

Then to $O(\lambda^6)$

$$V_{us} = \lambda, V_{cb} = A\lambda^2,$$

$$V_{ub} = A\lambda^3(\bar{\rho} - i\bar{\eta}),$$

$$V_{td} = A\lambda^3(1 - \bar{\rho} - i\bar{\eta}),$$

$$\text{Im}V_{cd} = -A\lambda^5\eta,$$

$$\text{Im}V_{ts} = -A\lambda^4\eta,$$

$$\bar{\rho} = \rho\left(1 - \frac{\lambda^2}{2}\right), \bar{\eta} = \eta\left(1 - \frac{\lambda^2}{2}\right)$$

May 17-25, 2005

Course at University of Barcelona

Peter Krizán, Ljubljana



Unitary relations

Rows and columns of the V matrix are orthogonal

Three examples: 1st+2nd, 2nd+3rd, 1st+3rd columns

$$V_{ud}V_{us}^* + V_{cd}V_{cs}^* + V_{td}V_{ts}^* = 0,$$

$$V_{us}V_{ub}^* + V_{cs}V_{cb}^* + V_{ts}V_{tb}^* = 0,$$

$$V_{ud}V_{ub}^* + V_{cd}V_{cb}^* + V_{td}V_{tb}^* = 0.$$

Geometrical representation: triangles in the complex plane.

May 17-25, 2005

Course at University of Barcelona

Peter Krizán, Ljubljana



Unitary triangles

(a)

$$V_{ud}V_{us}^* + V_{cd}V_{cs}^* + V_{td}V_{ts}^* = 0,$$

$$V_{us}V_{ub}^* + V_{cs}V_{cb}^* + V_{ts}V_{tb}^* = 0, \rightarrow$$

$$V_{ud}V_{ub}^* + V_{cd}V_{cb}^* + V_{td}V_{tb}^* = 0.$$

(b)

(c)

7-92 7204A4

All triangles have the same area $J/2$ (about 4×10^{-5})

$$J = c_{12}c_{23}c_{13}^2 s_{12}s_{23}s_{13} \sin \delta$$

May 17-25, 2005

Course at University of Barcelona


Peter Kržan, Ljubljana



Unitarity triangle

THE unitarity triangle:

$$V_{ud}V_{ub}^* + V_{cd}V_{cb}^* + V_{td}V_{tb}^* = 0$$




Another notation:

$$\phi_1 = \beta$$

$$\phi_2 = \alpha$$

$$\phi_3 = \gamma$$



May 17-25, 2005

Course at University of Barcelona

Peter Kržan, Ljubljana





Angles of the unitarity triangle

$$\alpha \equiv \phi_2 \equiv \arg\left(\frac{V_{td}V_{tb}^*}{V_{ud}V_{ub}^*}\right)$$

$$\beta \equiv \phi_1 \equiv \arg\left(\frac{V_{cd}V_{cb}^*}{V_{td}V_{tb}^*}\right)$$

$$\gamma \equiv \phi_3 \equiv \arg\left(\frac{V_{ud}V_{ub}^*}{V_{cd}V_{cb}^*}\right) \equiv \pi - \alpha - \beta$$



May 17-25, 2005



Course at University of Barcelona

Peter Krizan, Ljubljana




b decays

Tree



EW penguin



May 17-25, 2005

Course at University of Barcelona

Peter Krizan, Ljubljana



Decay amplitude structure

Quark diagrams: classified in tree, penguin and electroweak penguin contributions.

B decays are not so easy: due to long distance physics effects; rescattering processes can change the quark content of the final state.

Describe the weak-phase structure of B-decay amplitude: sum of three terms with definite CKM coefficients:

$$A(q\bar{q}q') = V_{tb}V_{tq'}^*P_{q'}^t + V_{cb}V_{cq'}^*(T_{c\bar{c}q'}\delta_{qc} + P_{q'}^c) + V_{ub}V_{uq'}^*(T_{u\bar{u}q'}\delta_{qu} + P_{q'}^u)$$

May 17-25, 2005

Course at University of Barcelona

Peter Kriz̃an, Ljubljana



Decay amplitude structure: qqs and qqd decays

Use the unitarity condition to simplify the expressions for individual amplitudes:

$$A(c\bar{c}s) = V_{cb}V_{cs}^*(T_{c\bar{c}s} + P_s^c - P_s^t) + V_{ub}V_{us}^*(P_s^u - P_s^t),$$

$$A(u\bar{u}s) = V_{cb}V_{cs}^*(P_s^c - P_s^t) + V_{ub}V_{us}^*(T_{u\bar{u}s} + P_s^u - P_s^t),$$

$$A(s\bar{s}s) = V_{cb}V_{cs}^*(P_s^c - P_s^t) + V_{ub}V_{us}^*(P_s^u - P_s^t).$$

Nice feature: penguin amplitudes only come as differences – only in this way they are meaningful.

$$A(c\bar{c}d) = V_{tb}V_{td}^*(P_d^t - P_d^u) + V_{cb}V_{cd}^*(T_{c\bar{c}d} + P_d^c - P_d^u),$$

$$A(u\bar{u}d) = V_{tb}V_{td}^*(P_d^t - P_d^c) + V_{ub}V_{ud}^*(T_{u\bar{u}d} + P_d^u - P_d^t),$$

$$A(s\bar{s}d) = V_{tb}V_{td}^*(P_d^t - P_d^u) + V_{cb}V_{cd}^*(P_d^c - P_d^u).$$

May 17-25, 2005

Course at University of Barcelona

Peter Kriz̃an, Ljubljana



Low-energy effective Hamiltonians

Low-energy effective Hamiltonians: constructed using the operator product expansion (OPE):

$$\langle f | \mathcal{H}_{\text{eff}} | i \rangle \propto \sum_k \langle f | Q_k(\mu) | i \rangle C_k(\mu)$$

μ is an appropriate renormalization scale $O(m_b)$. The OPE allows one to separate the "long-distance" contributions to that decay amplitude from the "short-distance" parts.

"long-distance" contributions not calculable \rightarrow nonperturbative hadronic matrix elements

"short-distance" described by perturbatively calculable Wilson coefficient functions $C_k(\mu)$.

For B decays:

$$\mathcal{H}_{\text{eff}}(\Delta B = -1) = \frac{G_F}{\sqrt{2}} \left[\sum_{j=u,c} V_{jq}^* V_{jb} \left\{ \sum_{k=1}^2 Q_k^{jq} C_k(\mu) + \sum_{k=3}^{10} Q_k^q C_k(\mu) \right\} \right]$$



Decay asymmetry predictions - overview

Five classes of B decays.

Classes 1 and 2 are expected to have relatively small direct CP violations \rightarrow particularly interesting for extracting CKM parameters from interference of decays with and without mixing.

In the remaining three classes, direct CP violations could be significant, decay asymmetries cannot be cleanly interpreted in terms of CKM phases.

1. Decays dominated by a single term: $b \rightarrow ccs$ and $b \rightarrow sss$. SM cleanly predicts zero (or very small) direct CP violations because the second term is Cabibbo suppressed. Any observation of large direct CP-violating effects in these cases would be a clue to beyond Standard Model physics. The modes $B^+ \rightarrow J/\psi K^+$ and $B^+ \rightarrow \phi K^+$ are examples of this class. The corresponding neutral modes have cleanly predicted relationships between CKM parameters and the measured asymmetry from interference between decays with and without mixing.



Decay asymmetry predictions - overview

2. Decays with a small second term: $b \rightarrow ccd$ and $b \rightarrow uud$. The expectation that penguin-only contributions are suppressed compared to tree contributions suggests that these modes will have small direct CP violation effects, and an approximate prediction for the relationship between measured asymmetries in neutral decays and CKM phases can be made.

3. Decays with a suppressed tree contribution: $b \rightarrow uus$. The tree amplitude is suppressed by small mixing angles, $V_{ub}V_{us}$. The no-tree term may be comparable or even dominate and give large interference effects. An example is $B \rightarrow \rho K$.

May 17-25, 2005

Course at University of Barcelona

Peter Krizan, Ljubljana



Decay asymmetry predictions - overview

4. Decays with no tree contribution: $b \rightarrow ssd$. Here the interference comes from penguin contributions with different charge $2/3$ quarks in the loop. An example is $B \rightarrow KK$.

5. Radiative decays: $b \rightarrow s\gamma$. The mechanism here is the same as in case 4 except that the leading contributions come from electromagnetic penguins. An example is $B \rightarrow K^*\gamma$.

May 17-25, 2005

Course at University of Barcelona

Peter Krizan, Ljubljana



Decay asymmetry predictions – overview

b->qq̄s

$B \rightarrow q\bar{q}s$ Decay Modes

Quark Process	Leading Term	Secondary Term	Sample B_d Modes	B_d Angle	Sample B_s Modes	B_s Angle
$b \rightarrow c\bar{c}s$	$V_{cb}V_{cs}^* = A\lambda^2$ tree + penguin (c-t)	$V_{ub}V_{us}^* = A\lambda^4(\rho - i\eta)$ penguin only (u-t)	$J/\psi K_S$	β	$\psi\eta'$ $D_s\bar{D}_s$	β_S
$b \rightarrow s\bar{s}s$	$V_{cb}V_{cs}^* = A\lambda^2$ penguin only (c-t)	$V_{ub}V_{us}^* = A\lambda^4(\rho - i\eta)$ penguin only (u-t)	ϕK_S	β	$\phi\eta'$	0
$b \rightarrow u\bar{u}s$	$V_{cb}V_{cs}^* = A\lambda^2$ penguin only (c-t)	$V_{ub}V_{us}^* = A\lambda^4(\rho - i\eta)$ tree + penguin (u-t)	$\pi^0 K_S$ ρK_S	competing terms	$\phi\pi^0$ $K_S K_S$	competing terms
$b \rightarrow c\bar{u}s$	$V_{cb}V_{cs}^* = A\lambda^3$	0	$D^0 K$ ↘ common $\bar{D}^0 K$ ↗ modes	γ	$D^0\phi$ ↘ common $\bar{D}^0\phi$ ↗ modes	γ

May 17-25, 2005

Course at University of Barcelona

Peter Kriz̄an, Ljubljana



Decay asymmetry predictions – overview

b->qq̄d

$b \rightarrow q\bar{q}d$ Decay Modes

Quark Process	Leading Term	Secondary Term	Sample B_d Modes	B_d Angle * (leading terms only)	Sample B_s Modes	B_s Angle * (leading term)
$b \rightarrow c\bar{c}d$	$V_{cb}V_{cd}^* = -A\lambda^3$ tree + penguin (c-u)	$V_{ub}V_{ud}^* = A\lambda^3(1 - \rho - i\eta)$ penguin only (t-u)	$D^+ D^-$	* β	ψK_S	β_S
$b \rightarrow s\bar{s}d$	$V_{cb}V_{cd}^* = A\lambda^3(1 - \rho - i\eta)$ penguin only (t-u)	$V_{ub}V_{ud}^* = A\lambda^3$ penguin only (c-u)	$\phi\pi$ $K_S\bar{K}_S$	competing terms	ϕK_S	competing terms
$b \rightarrow u\bar{u}d$	$V_{cb}V_{cd}^* = A\lambda^3(\rho - i\eta)$ tree + penguin (uc)	$V_{ub}V_{ud}^* = A\lambda^3(1 - \rho - i\eta)$ penguin only (t-c)	$\pi\pi; \pi\rho$ πa_1	* α	$\pi^0 K_S$ $\rho^0 K_S$	competing terms
$b \rightarrow c\bar{u}d$	$V_{cb}V_{cd}^* = A\lambda^2$	0	$D^0\pi^0$ ↘ common $\bar{D}^0\pi^0$ ↗ modes	γ	$D^0 K_S$ ↘ common $\bar{D}^0 K_S$ ↗ modes	γ


May 17-25, 2005

Course at University of Barcelona

Peter Kriz̄an, Ljubljana



Decay asymmetry predictions – example $\pi^+ \pi^-$



N.B.: for simplicity we have neglected possible penguin amplitudes (which we know is wrong), do it properly later.

May 17-25, 2005

Course at University of Barcelona

Peter Krizan, Ljubljana



Decay asymmetry predictions – example $J/\psi K_S$

$b \rightarrow c\bar{c}s$: tree + penguin contribution $\sim V_{cb} V_{cs}^* = A\lambda^2$
 penguin only contribution $\sim V_{ub} V_{us}^* = A\lambda^4(\rho - i\eta)$

Take into account that we measure the $\pi^+ \pi^-$ component of K_S – also need the $(q/p)_K$ for the K system

$$\lambda_{\psi K_S} = \eta_{\psi K_S} \left(\frac{V_{tb}^* V_{td}}{V_{tb} V_{td}^*} \right) \left(\frac{V_{cs}^* V_{cb}}{V_{cs} V_{cb}^*} \right) \left(\frac{V_{cd}^* V_{cs}}{V_{cd} V_{cs}^*} \right) =$$

$$= \eta_{\psi K_S} \left(\frac{V_{tb}^* V_{td}}{V_{tb} V_{td}^*} \right) \left(\frac{V_{cb} V_{cd}^*}{V_{cb}^* V_{cd}} \right)$$

$$\text{Im}(\lambda_{\psi K_S}) = \sin 2\beta$$

$$\beta \equiv \phi_1 \equiv \arg \left(\frac{V_{cd} V_{cb}^*}{V_{td} V_{tb}^*} \right)$$

May 17-25, 2005

Course at University of Barcelona

Peter Krizan, Ljubljana



b → c anti-c s CP=+1 and CP=-1 eigenstates

$$a_{f_{CP}} = -\text{Im}(\lambda_{f_{CP}}) \sin(\Delta mt)$$

Asymmetry sign depends on the CP parity of the final state f_{CP} , $\eta_{f_{CP}} = \pm 1$

$$\lambda_{f_{CP}} = \eta_{f_{CP}} \frac{q}{p} \frac{\bar{A}_{f_{CP}}}{A_{f_{CP}}}$$

J/ψ K_S (π⁺ π⁻): CP=-1

- J/ψ: P=-1, C=-1 (vector particle J^{PC}=1⁻⁻): CP=+1
- K_S (→ π⁺ π⁻): CP=+1, orbital ang. momentum of pions=0 →
P(π⁺ π⁻)=(π⁻ π⁺), C(π⁻ π⁺)=(π⁺ π⁻)
- orbital ang. momentum between J/ψ and K_S l=1, P=(-1)^l=-1

J/ψ K_L(3π): CP=+1

Opposite parity to J/ψ K_S (π⁺ π⁻), because K_L(3π) has CP=-1

May 17-25, 2005

Course at University of Barcelona

Peter Krizán, Ljubljana



The kaon case

The two K states have very different lifetimes

$$\tau_L = (5.17 \pm 0.04) \times 10^{-8} \text{ s}$$

$$\tau_S = (0.8927 \pm 0.009) \times 10^{-10} \text{ s}$$

The eigenstates are in this case defined by lifetimes

$$|K_S\rangle = p|K^0\rangle + q|\bar{K}^0\rangle$$

$$|K_L\rangle = p|K^0\rangle - q|\bar{K}^0\rangle$$

With the mass difference

$$\Delta m_K = m_L - m_S = (3.491 \pm 0.009) \times 10^{-15} \text{ GeV}$$

May 17-25, 2005

Course at University of Barcelona


Peter Krizán, Ljubljana



The kaon case

In this case

$$\Delta\Gamma_K \approx -2\Delta m_K$$



K^0 at $t=0$, evolution in time
Full line: K^0 , dotted: \bar{K}^0

T: in units of τ_s

After a few τ_s : left only K_L ,
roughly equal mixture of K^0
and \bar{K}^0

May 17-25, 2005

Course at University of Barcelona

Peter Krizan, Ljubljana



The kaon case

Define ϕ_{12} with

$$\frac{M_{12}}{\Gamma_{12}} = -\frac{|M_{12}|}{|\Gamma_{12}|} e^{i\phi_{12}}$$

It turns out that for the K system $\phi_{12} \ll 1$

From

(see above)

$$(\Delta m_B)^2 - \frac{1}{4}(\Delta\Gamma_B)^2 = 4(|M_{12}|^2 - \frac{1}{4}|\Gamma_{12}|^2)$$

$$\Delta m_B \Delta\Gamma_B = 4 \operatorname{Re}(M_{12} \Gamma_{12}^*)$$

To the leading order

$$\Delta\Gamma_K = -2|\Gamma_{12}|$$

$$\Delta m_K = 2|M_{12}|$$

May 17-25, 2005

Course at University of Barcelona

Peter Krizan, Ljubljana



Define

$$\Gamma_{12} = |\Gamma_{12}|e^{-2i\xi_K}$$

Use same
expression for q/p
as for the B case:

$$\frac{q}{p} = -\frac{\Delta m_B - \frac{i}{2}\Delta\Gamma_B}{2(M_{12} - \frac{i}{2}\Gamma_{12})} = -\frac{2(M_{12}^* - \frac{i}{2}\Gamma_{12}^*)}{\Delta m_B - \frac{i}{2}\Delta\Gamma_B}$$

May 17-25, 2005

Course at University of Barcelona

Peter Krizan, Ljubljana



The kaon case

$$\left(\frac{q}{p}\right)_K = e^{2i\xi_K} \left[1 - i\phi_{12} \frac{1 + i\frac{\Delta\Gamma_K}{2\Delta m_K}}{1 + \left(\frac{\Delta\Gamma_K}{2\Delta m_K}\right)^2} \right]$$

The ratio p/q is almost a pure phase (similar as in the B case)
-> CPV in mixing small in both cases (but for different
reasons: small lifetime diff in B, small phase in K system)

CPV in interference between mixing and decay:

$\lambda=1$ to $O(0.001)$ -> small

May 17-25, 2005

Course at University of Barcelona

Peter Krizan, Ljubljana



To next
order ->

$$\frac{q}{p} \frac{\bar{A}_{\pi\pi}}{A_{\pi\pi}} = 1 - i\phi_{12} \frac{1 + i\frac{\Delta\Gamma_K}{2\Delta m_K}}{1 + \left(\frac{\Delta\Gamma_K}{2\Delta m_K}\right)^2}$$

-> can be used to extract ϕ_{12}

But: it is not easy to transform from ϕ_{12} to electroweak parameters because of long distance (strong interaction) contribution M_{12} .

May 17-25, 2005

Course at University of Barcelona

Peter Krizan, Ljubljana



Backup slides

May 17-25, 2005

Course at University of Barcelona

Peter Krizan, Ljubljana



Parity of B^0

P: space inversion $P|B^0\rangle = -|B^0\rangle$

Why is the parity of B^0 (pseudoscalar meson) -1 ?

B^0 is composed of two quarks with spin $\frac{1}{2}$,
with total spin $J=0$.

The two quark spins are combined to $\frac{1}{2} \oplus \frac{1}{2} = 0$,
the relative angular momentum is $l=0$ (ground
bound state of \bar{b} in d).

Prostorski del valovne funkcije ima
parnost $(-1)^{l=0}=+1$.

Quark in antiquark have opposite parities
 \Rightarrow additional factor -1

May 17-25, 2005

Course at University of Barcelona

Peter Kržan, Ljubljana

$x' = \Lambda x$ transformation

$$i\gamma^\mu \frac{\partial \psi(x)}{\partial x^\mu} - m\psi(x) = 0 \quad \text{Dirac equation}$$

$$i\gamma^\mu \frac{\partial \psi'(x')}{\partial x'^\mu} - m\psi'(x') = 0$$

\Downarrow

$$\psi'(x') = S\psi(x) \quad \text{transformation for a bispinor}$$

$$S^{-1}\gamma^\mu S = \Lambda^\mu_\nu \gamma^\nu \quad \text{conclude}$$

$$\Lambda^\mu_\nu = \begin{pmatrix} 1 & & & \\ & -1 & & \\ & & -1 & \\ & & & -1 \end{pmatrix} \quad \text{space inversion}$$

$$S^{-1}\gamma^0 S = \gamma^0 \quad S^{-1}\gamma^k S = -\gamma^k$$

spinor za for
anti-particles ($E \leq 0$)

Transformation
of bispinor

compare

substitute

conclude

spinor for
particles ($E \geq 0$)

$$\psi'_{1,2} = S\psi_{1,2} = \psi_{1,2}$$

$$\psi'_{3,4} = S\psi_{3,4} = -\psi_{3,4}$$

May 17-25, 2005

Course at University of Barcelona

Peter Kržan, Ljubljana