



Univerza v Ljubljani

Recent results and future plans of the B factory Belle

Peter Križan

University of Ljubljana and J. Stefan Institute

December 16, 2005

University of Karlsruhe

Peter Križan, Ljubljana



Contents

Experimental apparatus: Belle at KEK-B

CP violation in the B system

Searching for New Physics: FCNC processes



- Observation of $b \rightarrow d$ penguins: $B \rightarrow \rho\gamma, \omega\gamma$ decays
- CP violation in $b \rightarrow s$ decays
- A_{fb} vs q^2 in $B \rightarrow K^* l^+ l^-$ decays



Plans for the future: a Super B factory



December 16, 2005

University of Karlsruhe

Peter Križan, Ljubljana










Three Angles: (ϕ_1, ϕ_2, ϕ_3) or (β, α, γ)



Big Questions: *Are determinations of angles consistent with determinations of the sides of the triangle ? Are angle determinations from loop and tree decays consistent ?*


December 16, 2005

University of Karlsruhe

Peter Križan, Ljubljana



Mixing Induced CP Violation



$$\rightarrow A_{CP}(t) = \frac{\Gamma(\bar{B}^0(t) \rightarrow f_{CP}) - \Gamma(B^0(t) \rightarrow f_{CP})}{\Gamma(\bar{B}^0(t) \rightarrow f_{CP}) + \Gamma(B^0(t) \rightarrow f_{CP})} = -\xi_f \sin 2\phi_1 \sin \Delta m_B t$$

$$\xi_f = \pm 1 \text{ for } CP = \pm 1$$


December 16, 2005

University of Karlsruhe

Peter Križan, Ljubljana



Principle of CPV Measurement



December 16, 2005

University of Karlsruhe

Peter Križan, Ljubljana



If there is **more than one diagram** and additional weak phases, there is the possibility of **direct CPV** and a new term with a $\cos(\Delta m \Delta t)$ time dependence.



$$P(B \rightarrow f_{CP}; \Delta t) = \frac{e^{-|\Delta t|/\tau_B}}{4\tau_B} [1 + q \cdot \{A \cos(\Delta m \Delta t) + S \sin(\Delta m \Delta t)\}]$$



with $q = \pm 1$

December 16, 2005

University of Karlsruhe

Peter Križan, Ljubljana







Evidence and Observation of Direct CP Violation in B Decays

DCPV in $B^0 \rightarrow \pi^+ \pi^-$ and $B^0 \rightarrow K^- \pi^+$,

hep-ex/0502035 (PRL 95, 101801(2005)); hep-ex/0507045

Asymmetries in the Dalitz plot of $B^\pm \rightarrow K^\pm \pi^+ \pi^-$

hep-ex/0509001

December 16, 2005

University of Karlsruhe

Peter Križan, Ljubljana



Importance of direct CPV in B decays

"The final, completely definitive death of any superweak theory will come from the observation of direct CP violation in the B system.....

Evidence for such direct CP violation would be given by the difference between the asymmetry parameters in a decay such as $B \rightarrow \pi^+ \pi^-$ from that of $B \rightarrow J/\psi K_S$. This can be considered the ϵ' experiment for the B system."

Lincoln Wolfenstein, 1999

[All CPV in the B system is due to a new superweak interaction. Expect no CPV in $\Delta B=1$ transitions.]

[e.g. J.M. Soares +L.Wolfenstein, PRD 47,421 1993.]

December 16, 2005

University of Karlsruhe

Peter Križan, Ljubljana



Direct CPV asymmetry in B Decays

Asymmetry in B decay rates

$$\begin{aligned}
 A_{dir} &\equiv \frac{\Gamma(\bar{B} \rightarrow \bar{f}) - \Gamma(B \rightarrow f)}{\Gamma(\bar{B} \rightarrow \bar{f}) + \Gamma(B \rightarrow f)} \\
 &= \frac{2r \sin \phi \sin \delta}{1 + r^2 + 2r \cos \phi \cos \delta} \\
 &\boxed{r = |P| / |T|, \phi = \text{weak phase diff}} \\
 &\quad \delta = \text{strong phase diff}
 \end{aligned}$$

The direct CP asymmetry (A_{dir}) can be large if two amplitudes have comparable sizes, different weak phases as well as a strong phase difference. This can happen in certain B decays due to the interference of penguin (P) and tree (T) decays.


December 16, 2005

University of Karlsruhe

Peter Križan, Ljubljana



Hierarchy of diagrams for $B \rightarrow K\pi, \pi\pi$ decays



Possibility of tree-penguin interference.

N.B. in $B \rightarrow \pi\pi$ the two diagrams are the same order in λ

December 16, 2005

University of Karlsruhe

Peter Križan, Ljubljana



Experimental Situation for $B \rightarrow \pi^+ \pi^-$ in 2004



Belle 152 M $\bar{B}B$

with 372 ± 32 $B^0 \rightarrow \pi^+ \pi^-$ events

$$S_{\pi\pi} = -1.00 \pm 0.21 \pm 0.07$$

$$A_{\pi\pi} = +0.58 \pm 0.15 \pm 0.07$$

PRL 93, 021601 (2004)

5.2σ CPV,

First evidence for DCPV (3.2σ)



BABAR 227 M $\bar{B}B$

with 467 ± 33 $B^0 \rightarrow \pi^+ \pi^-$ events

$$S_{\pi\pi} = -0.30 \pm 0.17 \pm 0.03$$

$$A_{\pi\pi} = +0.09 \pm 0.15 \pm 0.04$$

hep-ex/0501071, to appear in PRL

Also ~ 3.2 s discrepancy between Belle and BaBar


December 16, 2005

University of Karlsruhe

Peter Križan, Ljubljana



$B \rightarrow \pi^+ \pi^-$ time evolution



666 ± 43 $B \rightarrow \pi^+ \pi^-$ signal events

ΔE -Mbc 2D fits to individual time intervals

Bkg subtracted fit projections for $B \rightarrow \pi^+ \pi^-$

December 16, 2005

University of Karlsruhe

Peter Križan, Ljubljana



Summary of Belle $B^0 \rightarrow \pi^+ \pi^-$ CPV results

$$A_{\pi\pi} = +0.56 \pm 0.12 \pm 0.06$$

$$S_{\pi\pi} = -0.67 \pm 0.16 \pm 0.06$$

1st error statistical,
2nd systematic

- Compelling evidence for direct CP violation in $B \rightarrow \pi^+ \pi^-$ with 4.0σ significance
- Confirms previous Belle results.
- Isospin analysis for this mode alone gives (95.4% C.L) $0^\circ < \phi_2 < 19^\circ$ & $71^\circ < \phi_2 < 180^\circ$

December 10, 2005

University of Karlsruhe


Peter Križan, Ljubljana



Consistency Checks with time-integrated fits

$$A_{\pi\pi} = +0.52 \pm 0.14$$

Counting experiment consistent with unbinned time-dependent fits.







Visible indication of direct CP violation.

December 16, 2005

University of Karlsruhe

Peter Križan, Ljubljana









A new approach to direct CPV using the Dalitz plot in $B^\pm \rightarrow K^\pm \pi^+ \pi^-$ (hep-ex/0509001)

Sample used for $B^\pm \rightarrow K^\pm \pi^+ \pi^-$ study:

contains 2248 ± 79 B^- , 2038 ± 76 B^+

Fix the resonant substructure,
then allow both the phase and
amplitude to be different for B^+
and B^- decays.

For each resonant amplitude
replace $a e^{i\delta}$ with $a e^{i\delta} (1 \pm b e^{i\varphi})$



De

FIG. 9: Illustration of the amplitude parametrization with Eq. 7.

Peter Križan, Ljubljana



Evidence for CP Violation in the Decay $B^\pm \rightarrow \rho^0 K^\pm$

TABLE I: Results of the best fit to $K^\pm \pi^\pm \pi^\mp$ events in the B signal region. The first quoted error is statistical and the second is the model dependent uncertainty. The quoted A_{CP} significance is statistical only.

Channel	Fraction (%)	δ ($^\circ$)	b	φ ($^\circ$)	A_{CP} significance (σ)
$K^*(892)\pi^\pm$	$13.0 \pm 0.8^{+0.5}_{-0.7}$	0 (fixed)	$0.078 \pm 0.033^{+0.012}_{-0.003}$	$-18 \pm 44^{+5}_{-13}$	2.6
$K_0^*(1430)\pi^\pm$	$65.5 \pm 1.5^{+2.2}_{-1.9}$	$55 \pm 4^{+1}_{-5}$	$0.069 \pm 0.031^{+0.010}_{-0.008}$	$-123 \pm 16^{+4}_{-5}$	2.7
$\rho(770)^0 K^\pm$	$7.85 \pm 0.93^{+0.64}_{-0.59}$	$-21 \pm 14^{+14}_{-19}$	$0.28 \pm 0.11^{+0.07}_{-0.09}$	$-125 \pm 32^{+10}_{-85}$	3.9
$\omega(782)K^\pm$	$0.15 \pm 0.12^{+0.03}_{-0.02}$	$100 \pm 31^{+38}_{-21}$	0 (fixed)	—	—
$f_0(980)K^\pm$	$17.7 \pm 1.6^{+1.1}_{-3.3}$	$67 \pm 11^{+10}_{-11}$	$0.30 \pm 0.19^{+0.05}_{-0.10}$	$-82 \pm 8^{+2}_{-2}$	1.6
$f_2(1270)K^\pm$	$1.52 \pm 0.35^{+0.22}_{-0.37}$	$140 \pm 11^{+18}_{-7}$	$0.37 \pm 0.17^{+0.11}_{-0.04}$	$-24 \pm 29^{+14}_{-20}$	2.7
$f_X(1300)K^\pm$	$4.14 \pm 0.81^{+0.31}_{-0.30}$	$-141 \pm 10^{+8}_{-9}$	$0.12 \pm 0.17^{+0.04}_{-0.07}$	$-77 \pm 56^{+88}_{-43}$	1.0
Non-Res.	$34.0 \pm 2.2^{+2.1}_{-1.8}$	$\delta_1^{nr} = -11 \pm 5^{+3}_{-3}$ $\delta_2^{nr} = 185 \pm 20^{+62}_{-19}$	0 (fixed)	—	—
$\chi_{c0} K^\pm$	$1.12 \pm 0.12^{+0.24}_{-0.08}$	$-118 \pm 24^{+37}_{-38}$	$0.15 \pm 0.35^{+0.08}_{-0.07}$	$-77 \pm 94^{+154}_{-11}$	0.7



$$A_{CP}(B^\pm \rightarrow \rho^0 K^\pm) = 0.28 \pm 0.10^{+0.07}_{-0.09} \quad (3.9\sigma)$$

Significance varies from 3.7σ to 4.0σ depending on the model for the resonant substructure (add or remove modes, change nr model, cpv in $b \rightarrow u$ background).

December 16, 2005

University of Karlsruhe


Peter Križan, Ljubljana





Why FCNC decays?

Flavour changing neutral current (FCNC) processes (like $b \rightarrow s$, $b \rightarrow d$) are forbidden at the tree level in the Standard Model. Proceed only at low rate via higher-order loop diagrams. Ideal place to search for new physics.



December 16, 2005

University of Karlsruhe

Peter Križan, Ljubljana




A large number of $b \rightarrow s$ modes are known,
where are the $b \rightarrow d$ penguins ?

Suppressed by $|V_{tb}/V_{ts}|^2$ vs $b \rightarrow s\gamma$

Interesting:

Measurement of $|V_{tb}/V_{ts}|$

CP violation could be sizeable in SM (order 10%)



$$\frac{\mathcal{B}(B \rightarrow (\rho, \omega)\gamma)}{\mathcal{B}(B \rightarrow K^*\gamma)} = S_\rho \left| \frac{V_{td}}{V_{ts}} \right|^2 \left(\frac{1 - m_\rho^2/M_B^2}{1 - m_{K^*}^2/M_B^2} \right)^3 \zeta^2 [1 + \Delta R]$$

Addresses the same physics issue as B_s - B_s mixing (future Tevatron RunII +LHCb goal).

December 16, 2005

University of Karlsruhe

Peter Križan, Ljubljana




V_{td}/V_{ts} from $B \rightarrow \rho\gamma, \omega\gamma$

The measured branching fraction, $\mathcal{B}(B \rightarrow (\rho\omega)\gamma) = (1.34^{+0.34}_{-0.31} {}^{+0.14}_{-0.10}) \times 10^{-6}$, translates to

$$|V_{td}/V_{ts}| = 0.200^{+0.026}_{-0.025} (\text{exp.}) {}^{+0.038}_{-0.029} (\text{theo.}),$$

which is compatible with SM constraints based on fits using measurements of other CKM parameters.




Implications of Belle's observation of $b \rightarrow d \gamma$

Together with the evidence of $B \rightarrow K^0 \bar{K}$ modes, Belle has demonstrated the existence of a new quark level transition: $b \rightarrow d$

+ measurement of $|V_{td}/V_{ts}|$

$$\frac{\mathcal{B}(B \rightarrow (\rho/\omega)\gamma)}{\mathcal{B}(B \rightarrow K^*\gamma)} \propto \left| \frac{V_{td}}{V_{ts}} \right|^2$$

SU(3) breaking correction
weak annihilation diagram for $\mathcal{B}(B \rightarrow \rho/\omega\gamma)$



BELLE

How New Physics can contribute to $b \rightarrow s$

$B^0 \rightarrow \phi K^0$

"Internal Penguin"

New physics in loops?

$B^0 \rightarrow \eta' K^0$

Many new phases are possible in SUSY

December 16, 2005 University of Karlsruhe Peter Križan, Ljubljana

BELLE

Searching for new physics phases in CP violation measurements in $b \rightarrow s$ decays

Example:



no KM phase



$\mathbf{B} \rightarrow \mathbf{K}$ + $\mathbf{B} \rightarrow \mathbf{K}$



SM: $\sin 2\phi_1^{\text{eff}} = \sin 2\phi_1$ from $B \rightarrow J/\psi K^0$ ($b \rightarrow c \bar{c} s$)



unless there are other, non-SM particles in the loop

December 16, 2005 University of Karlsruhe Peter Križan, Ljubljana












Constraints on Wilson coefficients from $A_{FB}(B \rightarrow K^* l^- l^+)(q^2)$



Integrated FB asymmetry

$$A_{FB}(B \rightarrow K^* l^- l^+) = 0.50 \pm 0.12 \pm 0.02; (3.4\sigma)$$

control sample:

$$A_{FB}(B \rightarrow K^+ l^- l^+) = 0.10 \pm 0.14 \pm 0.01$$

Observed integrated A_{FB} rules out some radical New Physics Models with incorrect signs/magnitudes of C_9 and C_{10}

December 16, 2005


University of Karlsruhe

Peter Križan, Ljubljana




Results of the unbinned fit to q^2 and $\cos(\theta)$ distributions for ratios of Wilson coefficients.

A_{10}/A_7



$|A_7|$ constrained from $b \rightarrow s \gamma$ to be close to SM

A_9/A_7



Ref: hep-ex/0508009

	negative A_7	positive A_7
A_9/A_7	$-15.3^{+3.4}_{-4.8} \pm 1.1$	$-16.3^{+3.7}_{-5.7} \pm 1.4$
A_{10}/A_7	$10.3^{+5.2}_{-3.5} \pm 1.8$	$11.1^{+6.0}_{-3.9} \pm 2.4$



$$-1401 < A_9 A_{10} / A_7^2 < -26.4 \text{ at 95% C.L.}$$

Peter Križan, Ljubljana



$A_{FB}(B \rightarrow K^* l^+ l^-)[q^2]$ at Super B Factory

- Assume 1 year of running at $5 \times 10^{35} / \text{nb/sec}$
- $\rightarrow 5/\text{ab}$ integrated luminosity, 10 billion B mesons
- $\Delta A_9/A_9 \sim 11\%$, $\Delta A_{10}/A_{10} \sim 13\%$
- $- A_7$ fixed to SM value



December 16, 2005

University of Karlsruhe

Peter Križan, Ljubljana



Fundamental Questions in Flavor Physics

Are there New Physics Phases and New sources of CP Violation Beyond the SM ?

Experiments: $b \rightarrow s \text{ CPV}$, compare CPV angles from tree and loops

Are there new operators with quarks enhanced by New Physics ?

Experiments: $A_{FB}(B \rightarrow K^* l l)$, $B \rightarrow K \pi$ rates and asymmetries

Are there right-handed currents ?

Experiments: $b \rightarrow s \gamma \text{ CPV}$, $B \rightarrow V V$ triple-product asymmetries

Are there new flavor changing neutral currents ?

Experiments: $b \rightarrow s \nu \bar{\nu}$, D-Dbar mixing+CPV+rare, $\tau \rightarrow \mu \gamma$

These questions can only be answered at a Super B Factory.

December 16, 2005

University of Karlsruhe

Peter Križan, Ljubljana



Super B Factory Motivation

- Physics beyond the Standard Model (SM) must exist.
 - finite m_ν
 - gravity
 - If the LHC finds New Physics at the TeV scale,
 - its flavor structure must be examined experimentally. A super B factory is the best tool for this purpose.
- cf. Physics of top quark
First estimate of mass: BB mixing → ARGUS
Direct production, Mass, width etc. → CDF/D0
Off-diagonal couplings, phase → BaBar/Belle
- $b \bar{b} \rightarrow u c t \bar{t}$ matter
 $d \bar{d} \rightarrow \bar{u} \bar{c} \bar{t} \bar{t}$ no antimatter
- $V_{CKM} = \begin{pmatrix} V_{ud} & V_{us} & V_{ub} \\ V_{cd} & V_{cs} & V_{cb} \\ V_{td} & V_{ts} & V_{tb} \end{pmatrix}$
- If the LHC finds nothing but a SM-like Higgs,
 - searching for deviations from the SM in flavor physics will be one of the best ways to find new physics.


December 16, 2005

University of Karlsruhe

Peter Križan, Ljubljana



A Broad Unbiased Approach to New Physics




December 16, 2005

University of Karlsruhe

Peter Križan, Ljubljana



CPV in $b \rightarrow s$ and diagnosis of new physics



Many other examples of using correlations to distinguish new physics scenarios have been examined.

T.Goto, Y.Okada, Y.Shimizu,T.Shindou, M.Tanaka (2002, 2004) + SuperKEKB LoI

December 16, 2005


University of Karlsruhe

Peter Križan, Ljubljana





Comparison of Super-B and LHCb

- Clean environment → measurements that no other experiment can perform. Examples: CPV in $B \rightarrow \phi K^0$, $B \rightarrow \eta' K^0$ for new phases, $B \rightarrow K_{S\Gamma^0}\gamma$ for right-handed currents.
- "B-meson beam" technique → access to new decay modes.
Example: discover $B \rightarrow K\nu\nu$.
- Measure new types of asymmetries
Example: forward-backward asymmetry in $b \rightarrow s\mu\mu$, see
- Rich, broad physics program including B , τ and charm physics
Examples: searches for $\tau \rightarrow \mu\gamma$ and $D-D$ mixing with unprecedented sensitivity.
- No other experiment can compete for New Physics reach in the quark sector.



December 16, 2005

University of Karlsruhe



BELLE

Requirements for the Super B detector


Critical issues at $L = 4 \times 10^{35}/\text{cm}^2/\text{sec}$

- ▶ **Higher background ($\times 20$)**
 - radiation damage and occupancy
 - fake hits and pile-up noise in the EM
- ▶ **Higher event rate ($\times 10$)**
 - higher rate trigger, DAQ and computing
- ▶ **Require special features**
 - low p_μ identification $\leftarrow s\mu\mu$ recon. eff.
 - hermeticity $\leftarrow v$ "reconstruction"

Possible solution:

- ▶ Replace inner layers of the vertex detector with a silicon striplet detector.
- ▶ Replace inner part of the central tracker with a silicon strip detector.
- ▶ Better particle identification device
- ▶ Replace endcap calorimeter by pure CsI.
- ▶ Faster readout electronics and computing system.

December 16, 2005 University of Karlsruhe Peter Križan, Ljubljana



B
BELLE

PID upgrade in the endcap

improve K/ π separation in the forward (high mom.) region for few-body decays of B's
good K/ π separation for $b \rightarrow d\gamma$, $b \rightarrow s\gamma$
improve purity in fully reconstructed B decays
low momentum ($< 1\text{GeV}/c$) e/ μ / π separation ($B \rightarrow K\eta$)
keep high the efficiency for tagging kaons

December 16, 2005 University of Karlsruhe Peter Križan, Ljubljana

B
BELLE

Proximity focusing RICH in the forward region

aerogel photon detector

Cherenkov photons
charged particle

2 cm 20 cm

K/ π separation at $4\text{ GeV}/c$
 $\theta_c(\pi) \sim 308\text{ mrad}$ ($n = 1.05$)
 $\theta_c(\pi) - \theta_c(K) \sim 23\text{ mrad}$

$d\theta_c(\text{meas.}) = \sigma_0 \sim 13\text{ mrad}$
With 20mm thick aerogel and 6mm PMT pad size


→ 6σ separation with $N_{pe} \sim 10$

December 16, 2005 University of Karlsruhe Peter Križan, Ljubljana



Beam test: Cherenkov angle resolution and number of photons

Beam test results with 2cm thick aerogel tiles:
 $>4\sigma K/\pi$ separation



-> Number of photons has to be increased.

December 16, 2005

University of Karlsruhe


Peter Križan, Ljubljana



How to increase the number of photons?

What is the optimal radiator thickness?

Use beam test data on σ_0 and N_{pe}



Minimize the error per track:

$$\sigma_{track} = \sigma_0 / \sqrt{N_{pe}}$$

Optimum is close to 2 cm

December 16, 2005

University of Karlsru




Radiator with multiple refractive indices

How to increase the number of photons without degrading the resolution?

-> stack two tiles with different refractive indices: "focusing" configuration

normal



December 16, 2005




University of Karlsruhe

Peter Križan, Ljubljana






Focusing configuration – data

4cm aerogel single index



2+2cm aerogel




December 16, 2005



Tiling of the radiator

Minimize photon yield losses at the aerogel tile boundary: hexagonal tiling scheme



- Cut into hexagonal shape from a square block
- Machining device: use "water-jet" thanks to the hydrophobic nature


December 16, 2005

University of Karlsruhe

Peter Križan, Ljubljana



KEKB Collider Upgrade Scenario



December 16, 2005

University of Karlsruhe


Peter Križan, Ljubljana



Yamauchi's Schedule for Super B

A Super B proposal was submitted to MEXT in August 2005.

KEKB/Belle project receives a grade of S(i.e. A+) in gov. review
A search for a new KEK laboratory director is underway.



December 16, 2005

University of Karlsruhe


Peter Križan, Ljubljana



Yamauchi's budget for Super B

SuperKEKB 年次計画 (2005.1.11)

総額: 465.8 Oku-yen



December 16, 2005

University of Karlsruhe

Peter Križan, Ljubljana



Summary 1

- Observation of direct CP violation in $B^0 \rightarrow \pi^+\pi^-$ and $K^+\pi^-$ decays, evidence in $B^- \rightarrow \rho^0 K^-$
- CP violation in $b \rightarrow s$ transitions remains below SM expectation, but statistically limited.
- Forward-backward asymmetry (A_{FB}) in $b \rightarrow s l^+ l^-$ is becoming another powerful tool to search for physics beyond SM.
- We are entering an exciting phase of precision measurements.

.... and there are much more interesting results, but could not be covered in this talk!

December 16, 2005

University of Karlsruhe

Peter Križan, Ljubljana



Summary 2

- B factories have proven to be an excellent tool for flavour physics
- Reliable long term operation, constant improvement of the performance.
- Short term plan: increase luminosity $\times 3$ by a crab cavity
- Major upgrade in 2009-10 -> Super B factory, $L \times 30$
- Essentially a new project, all components have to be replaced, plans exist (LoI), nothing is frozen...
- Expect a new, exciting era of discoveries, complementary to LHC
- Do not miss the chance to be part of it...

December 16, 2005

University of Karlsruhe

Peter Križan, Ljubljana